


Neuron

Sensory Experience Engages Microglia to Shape Neural Connectivity through a Non-Phagocytic Mechanism

Graphical Abstract



Authors

Lucas Cheadle, Samuel A. Rivera, Jasper S. Phelps, ..., Linda C. Burkly, Wei-Chung Allen Lee, Michael E. Greenberg

Correspondence

michael_greenberg@hms.harvard.edu

In Brief

Sensory experience induces Fn14 expression in relay neurons and TWEAK expression in microglia to drive refinement of retinogeniculate connectivity. Microglial TWEAK signals through neuronal Fn14 to eliminate a subset of synapses proximal to TWEAK-expressing microglia, whereas Fn14 acts alone at other synapses to strengthen connectivity.

Highlights

- Experience induces Fn14 expression in neurons and TWEAK expression in microglia
- Fn14 increases the number of spine-associated synapses when not bound by TWEAK
- Microglial TWEAK signals through neuronal Fn14 to locally decrease synapse numbers
- Microglia-driven synapse loss occurs through a non-phagocytic mechanism

Article

Sensory Experience Engages Microglia to Shape Neural Connectivity through a Non-Phagocytic Mechanism

Lucas Cheadle,^{1,5} Samuel A. Rivera,¹ Jasper S. Phelps,^{1,2} Katelin A. Ennis,³ Beth Stevens,⁴ Linda C. Burkly,³ Wei-Chung Allen Lee,⁴ and Michael E. Greenberg^{1,6,*}

¹Department of Neurobiology, Harvard Medical School, 220 Longwood Avenue, Boston, MA 02115, USA

²Program in Neuroscience, Harvard Medical School, 220 Longwood Avenue, Boston, MA 02115, USA

³Research and Early Development, Biogen, 115 Broadway, Cambridge, MA 04142, USA

⁴Department of Neurology, F.M. Kirby Neurobiology Center, Boston Children's Hospital, 300 Longwood Avenue, Boston, MA 02115, USA

⁵Present address: Cold Spring Harbor Laboratory, 1 Bungtown Road, Cold Spring Harbor, NY 11724, USA

⁶Lead Contact

*Correspondence: michael_greenberg@hms.harvard.edu

<https://doi.org/10.1016/j.neuron.2020.08.002>

SUMMARY

Sensory experience remodels neural circuits in the early postnatal brain through mechanisms that remain to be elucidated. Applying a new method of ultrastructural analysis to the retinogeniculate circuit, we find that visual experience alters the number and structure of synapses between the retina and the thalamus. These changes require vision-dependent transcription of the receptor Fn14 in thalamic relay neurons and the induction of its ligand TWEAK in microglia. Fn14 functions to increase the number of bulbous spine-associated synapses at retinogeniculate connections, likely contributing to the strengthening of the circuit that occurs in response to visual experience. However, at retinogeniculate connections near TWEAK-expressing microglia, TWEAK signals via Fn14 to restrict the number of bulbous spines on relay neurons, leading to the elimination of a subset of connections. Thus, TWEAK and Fn14 represent an intercellular signaling axis through which microglia shape retinogeniculate connectivity in response to sensory experience.

INTRODUCTION

The connectivity of the mature brain is established through a convergence of intrinsic biological factors (nature) and environmental cues (nurture). Although early stages of neural circuit assembly are governed by genetic programs *in utero*, nascent circuits are extensively refined in response to sensory experience during postnatal brain development (Katz and Shatz, 1996; Wiesel and Hubel, 1963). This dynamic process of sensory-dependent (SD) refinement tunes the connectivity of a given neuron by determining which of its immature synaptic connections are strengthened and maintained and which connections are eliminated. Impairments in synaptic refinement contribute to neurodevelopmental disorders such as autism and schizophrenia, and aberrant re-activation of refinement in the mature brain may contribute to neurodegeneration (Feinberg, 1982–1983; Hammond et al., 2019). Although these observations underscore the importance of refinement for brain function and human health, the cellular and molecular mechanisms through which sensory experience refines developing circuits remain incompletely understood.

Because many cellular processes occur simultaneously in the early postnatal brain, it has been difficult to study synaptic refine-

ment in isolation. This challenge has been addressed in part by studies of the retinogeniculate pathway of the mouse, a visual circuit that undergoes a robust phase of refinement across the first postnatal month (Hooks and Chen, 2020). In this circuit, retinal ganglion cell (RGC) axons synapse onto the dendrites of excitatory relay neurons in the dorsal lateral geniculate nucleus (dLGN) of the thalamus, which relay visual information to the cortex (Figure 1A). These connections between RGCs and relay neurons are complex, with each presynaptic RGC and its postsynaptic relay neuron partner sharing multiple synapses. Retinogeniculate refinement has been shown by electrophysiology to entail strengthening of a few immature RGC inputs onto a given relay neuron and concurrent elimination of inputs that fail to strengthen (Chen and Regehr, 2000). These physiological changes occur at the same time that structural remodeling of retinal axons that synapse onto their targets in the dLGN is observed (Hong et al., 2014). Although these changes in connectivity are initially driven by sensory-independent brain activity between birth and post-natal day 20 (P20), sensory experience drives further refinement during a critical period between P20 and P30 (Hooks and Chen, 2006, 2008).

Because of a lack of molecular analyses of the late postnatal dLGN, there is currently an incomplete understanding of the

mechanisms that regulate the SD refinement of the retinogeniculate circuit. In contrast, several studies have shown that molecules commonly associated with the innate immune system, including major histocompatibility complex (MHC) class I proteins, coordinate phases of retinogeniculate refinement that occur prior to the onset of experience (Corriveau et al., 1998; Datwani et al., 2009). More recently, it was discovered that microglia, the resident immune cells of the brain, sculpt neural circuits through phagocytic engulfment of presynaptic inputs during this early sensory-independent phase of development (Paolicelli et al., 2011; Schafer et al., 2012; Stevens et al., 2007; Tremblay et al., 2010). However, it has been unclear whether the functions of immune-related molecules in brain development are restricted to early phases of refinement or are also engaged to coordinate SD refinement between P20 and P30.

To address this gap in knowledge, we previously characterized SD gene expression in the dLGN using single-cell transcriptomics, identifying the cell-surface cytokine receptor *Fn14* (fibroblast growth factor-inducible protein, 14 kDa) as the most highly induced molecule in relay neurons in response to visual stimulation. *Fn14*, a member of the tumor necrosis factor (TNF) receptor superfamily, binds the TNF family cytokine TWEAK (TNF-associated weak inducer of apoptosis) (Wiley and Winkles, 2003), which, as we show here, is induced by visual experience selectively in microglia. Electrophysiological analysis in an *Fn14* knockout (KO) mouse demonstrated that, in the absence of *Fn14*, the connections between RGCs and relay neurons fail to strengthen and that these weak connections between RGCs and relay neurons are not properly eliminated. Notably, this dual requirement of *Fn14* for RGC input strengthening and elimination is restricted to the SD phase of refinement as earlier phases of circuit development proceed normally in the absence of *Fn14* (Cheadle et al., 2018).

In the current study, we combine a new method of automated serial transmission electron microscopy (TEM) (Maniates-Selvin et al., 2020; Graham et al., 2019) and Golgi staining to profile structural features of synapses at the peak of SD refinement in the dLGN. We find that *Fn14* expressed by relay neurons promotes an increase in the number of bulbous relay neuron spines that form synapses with RGC axons. However, when microglia expressing TWEAK are near thalamic relay neuron dendritic spines, the *Fn14*-dependent increase in spine numbers is suppressed. This effect of TWEAK appears not to involve microglial engulfment of synapses but, instead, requires experience-dependent signaling within dLGN relay neurons. Thus, we identified a mechanism through which experience engages microglia to shape synaptic connectivity during refinement of the retinogeniculate circuit.

RESULTS

Analysis of Retinogeniculate Synapses at Nanometer Resolution


In wild-type (WT) mice, visual function is established in part through the strengthening of retinogeniculate connections between P20 and P27 in response to visual experience. One feature that is thought to inform the strength of a retinogeniculate connection is the number of synapses shared by a RGC and a

given relay neuron in the dLGN (Hamos et al., 1987). Electrophysiological analysis revealed that when *Fn14* function is disrupted, synaptic strengthening fails to occur, and it has been suggested that this failure may arise from a decrease in the number of synapses linking RGCs to relay neurons (Cheadle et al., 2018). To determine whether *Fn14* strengthens connectivity by increasing the number of synapses formed by RGC inputs, we carried out an ultrastructural analysis of individual retinogeniculate connections in the dLGNs of an *Fn14* KO mouse and a WT littermate at P27, a time point when sensory experience has been shown by electrophysiology to actively strengthen synapses via a *Fn14*-dependent mechanism (Cheadle et al., 2018). We employed a new semi-automated serial TEM method called GridTape to visualize and quantify retinogeniculate synapses at nanometer resolution. This method leverages semi-automated section imaging and alignment, allowing us to reconstruct synapses in large volumes of dLGN tissue from a WT and a *Fn14* KO mouse in parallel while blinded to condition (Figure S1).

Using serial TEM of ultrathin 40-nm sections, we imaged ~6 million μm^3 regions from the dLGNs of a *Fn14* KO mouse and a WT mouse and identified the following features: (1) presynaptic RGC boutons, axonal compartments containing neurotransmitter-filled vesicles that converge upon dendrites of the dLGN and are distinguished from other inputs based on their large size and the presence of pale mitochondria (Figure 1B); (2) dendrites, postsynaptic branches of relay neurons that contain the molecular machinery for processing incoming neurotransmission from retinal boutons; (3) dendritic spines, actin-rich protrusions that contain neurotransmitter receptors and receive synaptic input from RGCs; and (4) synapses, specific loci where docked neurotransmitter vesicles in the presynaptic bouton release glutamate onto postsynaptic specializations of the relay neuron dendrite or dendritic spine.

In the WT dLGN, we found that 60% of RGC boutons made contact with dendritic segments of relay neurons containing spines, whereas 40% of RGCs converged on smooth segments of dendrite. RGC boutons are relatively large, and the majority not only contact spines but envelope the spines, forming multiple synapses onto the dendritic shaft and the spines themselves (Figures 1B and 1C). The convergence of RGC inputs onto spines is unique compared with other sensory-processing brain regions such as the cortex, in which presynaptic inputs are smaller and do not typically envelop spines (Morgan et al., 2016; Rafols and Valverde, 1973). Here we show that these uniquely complex interactions between RGC inputs and spines make up a significant proportion of retinogeniculate connections.

Because spines have not yet been systematically characterized in the mouse dLGN, we quantified the morphologies of spines within RGC boutons. Although live imaging of spines in the cortex has shown spines to exist along a structural continuum (Berry and Nedivi, 2017; Holtmaat and Svoboda, 2009), analyses of spines by techniques such as Golgi staining and electron microscopy (EM) have focused on the distinct characteristics of spine morphology as viewed in fixed tissue. These characteristics include spine length and spine head diameter (Parajuli et al., 2017; Peters and Kaiserman-Abramof, 1970; Sorra and Harris, 2000). In this study, we measured these objective structural parameters as well as the numbers of spines






Figure 2. Developmental Changes in Spines Require Experience and Postsynaptic Fn14

(A) Low- and high-magnification images of a Golgi-stained brain section centered on the dLGN (outlined). Scale bars: 200 (Ai), 100 (Aii), and 10 (Aiii) μm .

(B) Schematic of spine types defined in the study.

(C) Example images of Golgi-stained spines in *Fn14^{fl/fl}*; *VGLUT2-Cre* mice, *Fn14^{fl/fl}*; *Chx10-Cre* mice, and Cre-negative littermates at P27. As in other panels, contrast was increased to better display spine morphology. Arrows, bulbous spines. Scale bars, 2 μm .

(D) Bulbous spine density is decreased by genetic ablation of Fn14 in postsynaptic relay neurons.

(E) Bulbous spine density is unaffected by genetic ablation of Fn14 in presynaptic RGCs.

(F) Spine length is decreased by genetic ablation of Fn14 in relay neurons.

(G) Spine length is decreased by genetic ablation of Fn14 in RGCs.

(legend continued on next page)

corresponding to structurally defined subclasses: “bulbous spines” (similar to “mushroom” spines in other systems), which we defined as having a relatively large spine head that is at least twice as wide as the spine neck; “thin spines,” defined by a uniform diameter (head-to-neck width ratio less than 2) of less than 0.5 μm ; and spines of uniform diameter greater than 0.5 μm , which we called “broad non-bulbous spines” (these correspond in part to “stubby spines” in other systems) (Figure S2A). The results of hierarchical clustering of the distribution of spines based on the head-to-neck width ratio were consistent with these morphological categories representing distinct classes (Figures S2B and S2C).

Applying these classifications to our WT dataset, we found that the majority of spines that were enveloped by retinal inputs, about 60%, have a bulbous morphology. Although retinal inputs represent only about 10% of total synaptic inputs to the dLGN, synaptic contacts deriving from other brain regions, most prominently the visual cortex, were observed to converge on bulbous spines less frequently, suggesting that bulbous spines may be specialized to mediate incoming sensory information from the retina. Interestingly, we found that the different classes of spines contained varying numbers of synapses: bulbous spines contained, on average, ~ 4 synapses per spine, non-bulbous spines ~ 2 synapses per spine, and thin spines ~ 1 synapse per spine, suggesting that the different spine types make distinct contributions to synaptic strength. Based on this observation, bulbous spines, which contain the most synapses, are likely to be mediators of strong retinogeniculate connections.

Fn14 Increases Retinal Bouton Convergence onto Bulbous Spines

Analysis of retinogeniculate synapses across the dLGN of a P27 Fn14 KO mouse revealed that 30% of RGC inputs converged on spines in the absence of Fn14, in contrast to 60% in WT mice (Figures 1B–1D). Classification of spines by morphological subtype in the Fn14 KO dLGN revealed that this decrease in spine-convergent inputs is selective for RGC inputs converging on bulbous spines (Figure 1E). Not only was the number of retinal input-contacted bulbous spines $\sim 50\%$ lower in the Fn14 KO mouse (Figure 1F), but the bulbous spines that were associated with retinal inputs contained $\sim 40\%$ fewer synapses on average (Figure 1G). It is not clear whether the decrease in bulbous spines overall and the decrease in synapses per spine represent distinct parallel mechanisms or are two steps of a single process. In either case, the decrease in bulbous spine-associated synapses contributed to a substantial 30% decrease in the total number of

synapses connecting the retina to the dLGN in the absence of Fn14. In contrast, the numbers of synapses associated with dendritic shafts, non-bulbous spines, and thin spines in the Fn14 KO dLGN were equivalent to those in the WT (Figures 1H and S2D–S2I), suggesting that the regulation of synapses by Fn14 is selective for bulbous spines. Given that the reported deficits in synaptic strengthening in the Fn14 KO correlate with a significant decrease in bulbous spine-associated synapses, we reasoned that the formation or maintenance of synapse-containing bulbous spines by Fn14 is likely to underlie SD synaptic strengthening in the dLGN.

Developmental Spine Changes Are Coordinated by Experience and Postsynaptic Fn14

We next employed morphological spine analysis via Golgi staining as a surrogate readout of the changes in synaptic strength that occur during postnatal dLGN development. We performed Golgi staining on the dLGNs of constitutive Fn14 KO and WT mice at P20 and P27, the time points flanking the SD phase of retinogeniculate refinement, to determine whether we can detect the decrease in bulbous spines we had observed by TEM using this more tractable method. We found that spines were normal in the absence of Fn14 at P20 (Figure 1I) but that the number of bulbous spines was decreased significantly in the Fn14 KO dLGN compared with the WT at P27 (Figure 1J), consistent with our TEM results (Figure 1F).

Because the analyses described so far were performed in a constitutive KO mouse lacking Fn14 in all cell types, it remained unclear whether Fn14 regulates synaptic refinement through a presynaptic mechanism, a postsynaptic mechanism, or both. To address this gap in knowledge, we developed a Fn14^{fl/fl} conditional KO mouse in which exons 2–4 of the *Fn14* locus are flanked by LoxP sites and compared spine numbers in Fn14^{fl/fl} Cre-negative control mice with spine numbers in Fn14^{fl/fl} mice crossed to a relay neuron-specific VGLUT2-Cre driver or a retina-specific Chx10-Cre driver (Rowan and Cepko, 2004). We validated region-specific loss of Fn14 in Cre-positive mice by protein and mRNA analysis (Figures S3A and S3B). We found that Fn14 expression in relay neurons is required to form or maintain bulbous spines, consistent with Fn14 functioning to regulate synapse number and strength via a postsynaptic mechanism that is consistent with its SD transcription in relay neurons (Figures 2A–2D). In contrast, genetic ablation of Fn14 in RGCs had no significant effect on bulbous spine numbers in the dLGN (Figure 2E). However, we found that spine length is significantly decreased by ablation of Fn14 in pre- or postsynaptic neurons,

(H) Examples of Golgi-stained dendrites and spines, analyzed across postnatal development and in animals subjected to late dark rearing (LDR). Arrows, bulbous spines. Scale bar, 2 μm .

(I) Bulbous spine density increases between P20 and P27 in WT mice.

(J) Bulbous spine density is decreased significantly in LDR mice compared with normally reared (NR) mice.

(K) Spine head diameter across all spine types is decreased by LDR.

(L) Thin spine density decreases between P10 and P20 in WT mice.

(M) Non-bulbous spine density increases between P10 and P20 and then decreases between P20 and P27 in WT mice.

(N) Spine length across all spine types increases during development.

(O) Spine head diameter increases between P10 and P27 in WT mice.

Statistical analysis: (D)–(G), (J), and (K), Student's t test; (I) and (L)–(O), one-way ANOVA with Tukey's post hoc comparison. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. Means are plotted with individual data points \pm SEM.




Figure 3. Experience Drives Expression of TWEAK in Microglia and Fn14 in Relay Neurons

(A) Single-cell RNA sequencing analysis of *TWEAK* expression across postnatal development in the dLGN. y axis, *TWEAK* mRNA transcripts per cell; x axis, age. (B) *TWEAK* protein levels, as measured by ELISA, increase across postnatal development. Levels are normalized to P12. (C) Confocal images of mRNA expression in dLGNs of mice subjected to LDR (unstimulated controls) or mice subjected to LDR and then acutely re-exposed to light for 8 h (+ light). Sections were probed for markers of microglia (*Cx3cr1*) or relay neurons (*VGLUT2*) (white), *TWEAK* (green), and *Fn14* (red). Scale bar, 5 μ m. (D) The percentage of microglia expressing at least 3 *TWEAK* mRNA molecules is increased significantly in light-stimulated mice. (E) Cumulative frequency distribution plot displaying increased *TWEAK* expression per microglia following light re-exposure. (F) Cumulative frequency distribution plot displaying increased *Fn14* expression per relay neuron following light re-exposure. (G) qPCR quantification of *TWEAK* expression in microglia isolated from visual cortices of mice following LDR and light re-exposure for 8–12 h. *TWEAK* expression was normalized to *Cx3cr1* expression to account for variability in enrichment efficiency. Statistical analysis: (A), (B), and (G), one-way ANOVA with Tukey's post hoc comparison; (D), Student's t test; (E) and (F), Kolmogorov-Smirnov distribution test. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. Means are plotted with individual data points \pm SEM.


highlighting that the functions of *Fn14* at synapses are likely to be multi-faceted (Figures 2F and 2G).

An analysis of spine morphology across postnatal development in WT mice revealed that bulbous spine numbers increase between P20 and P27, highlighting that visual experience likely engages *Fn14* during this period to regulate the number of bulbous spines during the course of development (Figures 2H and 2I). If so, then we would expect depriving mice of experience during this phase, a manipulation that results in a significant decrease in *Fn14* expression, to result in a decrease in the number of bulbous spines, phenocopying the *Fn14* KO mouse. We find that, compared with normally reared (NR) mice, mice reared in complete darkness between P20 and P27 (late dark rearing [LDR]) have 50% fewer bulbous spines, with the average diameter of all spines decreasing by 12% (Figures 2J and 2K).

Although changes in the overall structure of spines as well as in the numbers of thin and non-bulbous spines also occurred during postnatal development (Figures 2L–2O), the numbers of thin and non-bulbous spines and spine length overall were unaffected by LDR (Figures S2J–S2L). Altogether, our data suggest that *Fn14*-dependent addition of bulbous-spine-associated synapses to retinogeniculate connections is one mechanism by which experience strengthens connectivity during the vision-sensitive phase of retinogeniculate circuit maturation.

Sensory Experience Induces TWEAK Expression in Microglia

We next sought to find out whether the cytokine ligand of *Fn14*, *TWEAK*, collaborates with *Fn14* to promote neural circuit development. We first assessed the pattern of *TWEAK* expression by



(legend on next page)

interrogating a single-cell RNA sequencing dataset from the dLGN (Kalish et al., 2018). Although TWEAK is lowly expressed in multiple cell types at P5 and P10, TWEAK expression is significantly upregulated in microglia between P10 and P16, time points flanking the onset of visual experience at eye opening (Figures 3A and 3B). TWEAK expression continues to increase in microglia into the period of SD refinement that begins by P20.

The temporal overlap between onset of vision and upregulation of TWEAK in microglia led us to hypothesize that, in addition to its ability to induce gene expression in relay neurons, experience might also promote transcriptional changes in dLGN microglia. To test this idea, mice were subjected to LDR and then re-exposed to light for 8 h, a treatment that leads to Fn14 upregulation in relay neurons. Multiplexed single-molecule fluorescence *in situ* hybridization (smFISH) was then used to probe dLGN sections for *TWEAK*, *Fn14*, and cell-type-specific markers of microglia (*Cx3cr1*) or relay neurons (*VGLUT2*). We found that about 70% of microglia express *TWEAK* following visual stimulation, whereas 8% of dLGN microglia express *TWEAK* in unstimulated mice (Figures 3C and 3D). Analysis of the number of *TWEAK* mRNA molecules per microglia and the number of *Fn14* molecules per relay neuron revealed strong light-driven increases in *TWEAK* and *Fn14* expression in individual cells of non-overlapping classes (Figures 3E and 3F). A light-dependent increase in *TWEAK* expression was also observed by qPCR following isolation of microglia from the visual cortices of stimulated and unstimulated mice (Figure 3G). These patterns of stimulus-dependent *TWEAK* mRNA expression are largely recapitulated at the level of the TWEAK protein, as measured by ELISA (Figures S4A and S4B). Although we also detected low levels of TWEAK in astrocytes and endothelial cells (Figures 3A and S4C), TWEAK expression is notably higher in microglia, and only microglia induce TWEAK expression in response to visual experience (Figures S4D–S4F).

TWEAK Is Dispensable for Phagocytic Engulfment of Synapses by Microglia

We next considered what the function of TWEAK might be during retinogeniculate circuit refinement. Phagocytic engulfment of synapses is the best-characterized mechanism by which microglia shape developing brain circuits (Cowan and Petri, 2018; Ne-

niskyte and Gross, 2017). We therefore hypothesized that TWEAK expressed by microglia might mediate elimination of retinogeniculate synapses by binding to Fn14 expressed by relay neuron synapses, leading to their engulfment. This possibility is consistent with our previous finding that, in addition to a failure to strengthen RGC inputs to relay neurons in Fn14 KO mice, we also observed a failure to eliminate retinogeniculate connections that do not strengthen in response to visual experience.

When we assessed the degree of synaptic engulfment by microglia in the dLGNs of WT mice, we found that pre- but not post-synaptic elements are engulfed by microglia at the height of SD refinement (P27; Figure S5). When we compared pre-synaptic engulfment by microglia in WT and TWEAK KO mice (Dohi et al., 2009), we found that TWEAK is not required for phagocytic engulfment of synapses by microglia at an early time point that precedes visual experience (P7) or at the height of SD refinement (P27) (Figures 4A and 4B). Moreover, eye-specific segregation of ipsi- and contralateral retinal inputs, a developmental process that relies on synaptic engulfment by microglia (Schafer et al., 2012; Stevens et al., 2007), proceeds normally in TWEAK KO mice (Figures 4C and 4D). In addition, other aspects of microglial health and function, such as the number of microglia and their morphology, are also unaffected by disruption of TWEAK function (Figures 4E–4G). Together, these data suggest a possible function for SD TWEAK activation in microglia that does not involve engulfment of synapses.

Microglial TWEAK-Dependent Regulation of Bulbous Spines Requires Light and Postsynaptic Fn14

Having established spine analysis as a proxy for functional circuit changes downstream of experience and Fn14, we next assessed the effect of disrupting TWEAK function on the number and morphology of spines. Given that Fn14 promotes an increase in the number of bulbous spines in the dLGN, we hypothesized that, if TWEAK binding activates Fn14 function, then genetic ablation of TWEAK might lead to a decrease in the number of bulbous spines. Alternatively, if TWEAK binding to Fn14 inhibits Fn14 function, then disruption of TWEAK function might lead to an increase in the number of bulbous spines. We found that, in the absence of TWEAK, relay neurons display a significant increase in the number of bulbous spines. This

Figure 4. TWEAK Promotes Experience-Dependent Spine Loss through a Non-phagocytic Mechanism

- (A) Surface renderings of microglia (green) containing phagocytosed retinal boutons (blue and red) labeled by ocular injection of fluorophore-conjugated tracers in mice at P27. Microglial reconstructions are based on a combination of Iba1 and P2ry12 marker immunostaining. Scale bars, 10 μm (inset, 1 μm).
- (B) Quantification of the volume of individual microglia occupied by retinal inputs, plotted as the percentage of a given occupied microglial cell.
- (C) Confocal images of fluorescently labeled RGC boutons in contralateral (green) and ipsilateral (red) dLGNs. Microglia immunostained for Iba-1 are shown in white. Scale bars, 400 μm .
- (D) Quantification of the overlap between ipsi- and contralateral inputs, measured by co-localized signal in Imaris.
- (E) Example tracings of microglia analyzed by Sholl morphological analysis. Scale bars, 10 μm .
- (F) Sholl analysis of morphological complexity, indicating the number of microglial projections intersecting with a series of concentric circles radiating out from the soma.
- (G) Quantification of the number of microglia per dLGN volume based on Iba-1 staining.
- (H) Schematic of the viral overexpression experiments probing the role of TWEAK in spine development. Inset, FISH probing for *TWEAK* mRNA (green). Scale bars, 12 μm .
- (I) Example images of dendritic spines in the dLGNs of TWEAK KO and WT mice of different genotypes following viral infection. NR, normally reared. LDR, late dark reared. Arrows, bulbous spines. Scale bar, 2 μm .
- (J) Quantification of bulbous spine densities in TWEAK KO and WT mice following viral infection with or without experience. Con., control virus. Twe., TWEAK virus.

Statistical analysis: one-way ANOVA with Tukey's post hoc comparison. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$. Means are plotted with individual data points \pm SEM.




Figure 5. Microglial TWEAK Signals through Postsynaptic Fn14 to Decrease Bulbous Spine Numbers

(A) Example images of spines in the dLGNs of Fn14^{fl/fl} Cre-negative or VGLUT2-Cre-positive mice following viral infection. Scale bar, 2 μ m.
 (B) Quantification of bulbous spine densities following TWEAK or mCherry expression in the dLGNs of Cre-negative (–) and VGLUT2-Cre-positive (+) mice. A comparison of mCherry-infected conditions is also plotted in Figure 2D.
 (C) Example images of spines in the dLGNs of TWEAK^{fl/fl} Cre-negative or Cx3cr1-Cre-positive mice. Scale bar, 2 μ m.
 (D) Bulbous spine density is increased by genetic ablation of TWEAK in microglia.
 (E) Spine head diameter is increased by genetic ablation of TWEAK in microglia.
 (F) Thin spine density is unaffected by genetic ablation of TWEAK in microglia.
 (G) Non-bulbous spine density is unaffected by genetic ablation of TWEAK in microglia.
 (H) Total spine density is unaffected by genetic ablation of TWEAK in microglia.
 (I) Spine length is unaffected by genetic ablation of TWEAK in microglia.
 (J) Confocal images of dLGNs from a TWEAK KO mouse, a Fn14^{fl/fl} Cre-negative mouse, and a Fn14^{fl/fl}; VGLUT2-Cre+ mouse following bath application of recombinant mouse TWEAK and subsequent immunostaining for TWEAK (red) and VGLUT2 (green). Scale bar, 10 μ m.
 (K) Western blot of whole mouse forebrain fractionated to enrich for synaptosomes. Blots were probed for Fn14, the retinal presynaptic marker VGLUT2, the postsynaptic marker PSD-95, and GAPDH (a non-synaptic control).

(legend continued on next page)

suggests that TWEAK binding to Fn14 antagonizes the ability of Fn14 to enhance bulbous spine numbers or might initiate active disassembly of bulbous spines (Figures 4H–4J).

To further examine whether TWEAK antagonizes the effect of Fn14 on bulbous spine numbers, we overexpressed TWEAK in the dLGN by bilaterally injecting adeno-associated viruses (AAVs) expressing soluble TWEAK or mCherry (control) into the right and left dLGN, respectively, of TWEAK KO and WT mice at P12 (Figure 4H). We confirmed that, at the area of injection, glial activation is minimal in most of the dLGN sections analyzed 15 days after injection (Figure S6). The AAVs we injected predominantly led to soluble TWEAK expression in relay neurons of the dLGN (Figure 4H). Therefore, although this experiment does not allow us to specifically assess the role of microglial TWEAK in spine development, it nevertheless allowed us to determine whether localized overexpression of soluble TWEAK is sufficient to affect dLGN synapses *in vivo*. We assessed synaptic morphology in TWEAK-overexpressing mice by Golgi staining at P27. In a WT background, we found that overexpression of TWEAK was sufficient to decrease the number of bulbous spines by 25%. In contrast, mice in which TWEAK function was disrupted had 30% more bulbous spines than their WT littermates, and this increase in bulbous spines in KO mice was reversed by re-expression of TWEAK (Figures 4I and 4J).

Given that Fn14-dependent enhancement of synaptic strengthening and bulbous spine numbers are dependent on visual experience (Cheadle et al., 2018), we next wanted to find out whether the antagonistic effect of TWEAK on bulbous spine numbers also depends on sensory input. To test this possibility, we overexpressed soluble TWEAK in the dLGNs of WT mice at P12, subjected the mice to LDR between P20 and P27, and analyzed spines. Although we again found that LDR of WT mice leads to a significant decrease in the number of bulbous spines compared with WT mice housed under standard conditions (Figure 2J), TWEAK overexpression had no further effect on bulbous spine numbers in LDR mice, indicating that TWEAK signaling requires experience to decrease bulbous spine numbers (Figures 4I and 4J).

To test directly whether Fn14 expression in relay neurons is required for the TWEAK-dependent decrease in bulbous spines, we virally overexpressed soluble TWEAK or mCherry in Fn14^{fl/fl}; VGLUT2-Cre mice or Cre-negative littermates and assessed spines. As shown in Figure 2D, removal of Fn14 from relay neurons by crossing conditional KO mice to the VGLUT2-Cre line decreased bulbous spines by about 50%. We found that overexpressed TWEAK significantly decreased bulbous spine numbers in Cre-negative mice, in which relay neurons express normal levels of Fn14, but did not affect spine numbers in Fn14^{fl/fl}; VGLUT2-Cre mice, where Fn14 expression is selectively ablated in relay neurons (Figures 5A and 5B). This finding, taken together with the observation that TWEAK overexpression in the dLGNs of visually deprived mice has no effect on bulbous spine numbers (Figure 4J), indicates that the TWEAK-dependent

decrease in spines requires sensory-evoked Fn14 expression in relay neurons. Because we have shown that bulbous spines contain approximately four synapses per spine and interact with ~40% of incoming retinal terminals, these TWEAK-driven changes in bulbous spine numbers are likely to have a powerful functional effect on retinogeniculate connectivity.

SD regulation of spine numbers by TWEAK-Fn14 signaling may be a mechanism by which microglia drive synapse loss. However, the experiments presented so far using TWEAK KO mice, in which TWEAK function is disrupted in all cells, do not explicitly show that microglia are the relevant expressers of TWEAK in the context of synaptic development, especially given the observation that, in addition to microglia, endothelial cells and astrocytes express low levels of TWEAK (Figure S4C). Therefore, we next wanted to find out whether microglia-expressed TWEAK mediates the decrease in bulbous spine numbers. Toward this end, we generated a TWEAK floxed conditional KO mouse and crossed it to the Cx3cr1-Cre microglial driver (Yona et al., 2013; validation in Figures S3C–S3G). Analysis of dendritic spines in these mice at P27 revealed a 70% increase in the number of bulbous spines compared with WT mice (Figures 5C and 5D). We also noted a small but significant increase in spine head diameter upon microglial ablation of TWEAK, consistent with bulbous spines having relatively large head diameters (Figure 5E). Other spine parameters were unaffected by loss of microglial TWEAK (Figures 5F–5I). Together, these data identify a role of microglial TWEAK-to-neuronal Fn14 signaling in restricting the number of bulbous spines in response to sensory experience via a mechanism that does not involve synaptic engulfment.

TWEAK and Fn14 Are Likely to Signal Locally at Retinogeniculate Synapses

To further investigate the possibility that TWEAK and Fn14 signal locally at a subset of synaptic connections to regulate bulbous spine numbers, we developed a method for assessing the subcellular localization of TWEAK. Acute dLGN slices from a Fn14^{fl/fl}; VGLUT2-Cre or an Fn14^{fl/fl}; Cre-negative littermate were bathed with recombinant mouse TWEAK and then washed and immunostained for VGLUT2, a retinogeniculate synapse marker, and TWEAK using an antibody capable of detecting TWEAK when overexpressed (Chicheportiche et al., 1997). Strikingly, we found that, in mice expressing neuronal Fn14, recombinant TWEAK was highly localized to retinogeniculate synapses, whereas very little recombinant TWEAK was detected at synapses in the dLGN of mice lacking Fn14 in relay neurons, suggesting that recombinant TWEAK binds to synaptically localized Fn14 (Figure 5J). Consistent with Fn14-dependent localization of TWEAK at retinogeniculate synapses, synaptosomal fractionation of the mouse brain revealed enrichment of Fn14 at synapses (Figure 5K). In addition, an unbiased analysis of candidate Fn14 binding partners by co-immunoprecipitation and mass spectrometry identified a cohort of 24 likely Fn14 interactors

(L) Functional protein association network determined by STRING analysis, illustrating known and predicted interactions between proteins identified as potential Fn14 interactors by mass spectrometry.

Statistical analysis: (B), one-way ANOVA with Tukey's post hoc comparison; (D)–(I), Student's t test. **p* < 0.05, ***p* < 0.01, ****p* < 0.001. Means are plotted with individual data points ± SEM.




Figure 6. Microglial TWEAK Selectively Eliminates Proximal Bulbous Spines In Vitro

(A) Schematic of the *in vitro* co-culture experiment. Neurons from embryonic thalami (thal.) were sparsely transfected to express mCherry and then seeded with microglia isolated from TWEAK KO; Cx3cr1-GFP or WT; Cx3cr1-GFP mice. Spines were analyzed 24 h later.

(B) Confocal images of neurons (red) co-cultured with TWEAK KO or WT microglia (green). (Bi) mCherry-filled neuron contacted by WT GFP+ microglia. Scale bar, 15 μ m. (Bii) Example of WT microglia contacting the dendrite of a neuron. Scale bar, 5 μ m. (Biii) Example of TWEAK KO microglia contacting the dendrite of a neuron. Scale bar, 5 μ m. Shown are dendrites with spines from each condition; scale bar, 2 μ m.

(C) Total spine density in cultures without microglia (control) or with TWEAK WT or TWEAK KO microglia. Neurons co-cultured with TWEAK KO microglia maintained significantly more spines than those co-cultured with WT microglia.

(D) Quantification of bulbous spine density reveals that bulbous spines are protected when microglial TWEAK is ablated.

(E) Quantification of non-bulbous spine density in co-cultures.

(F) Quantification of thin spine density in co-cultures.

(G) Cumulative frequency distribution plot reflecting the proximity of bulbous spines to the nearest microglia with or without TWEAK expression. Bulbous spines were maintained closer to microglia when microglia lacked TWEAK.

(H) Violin plot reflecting median (dashed line) and quartile (dotted lines) values of the distance between bulbous spines and microglia with or without TWEAK.

(I) Cumulative frequency distribution plot reflecting the proximity of thin spines to the nearest microglia in co-cultures with WT microglia (black) or TWEAK KO microglia (purple).

(J) Violin plot reflecting median (dashed line) and quartile values (dotted lines) of the distance between thin spines and microglia with or without TWEAK.

Means are plotted with individual data points \pm SEM. Statistical analysis (C)–(F): one-way ANOVA with Tukey’s post hoc comparison; (G) and (I), Kolmogorov-Smirnov distribution comparison; (H) and (J), Student’s t test. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.




Figure 7. Synapse Number Is Negatively Correlated with the Level of TWEAK Expression in Nearby Microglia

(A) Stimulated emission depletion (STED) microscopy images of a microglial cell process, visualized by immunostaining for Iba1 and P2ry12 (green) within nanometer distance of VGLUT2-immunostained retinogeniculate synapses (red). Scale bars, 4 μ m; inset, 2 μ m.

(B) Confocal images and volumetric reconstructions of microglia (Iba1 and P2ry12, *Bi* and *Bii*, green), TWEAK mRNA (*Biii* and *Biv*, red), and retinogeniculate synapses (VGLUT2, *Bv* and *Bvi*, blue). (*Bvii*) and (*Bviii*) Channels merged. Arrows mark TWEAK mRNA transcripts within microglia. Top rows: microglia expressing only 1 mRNA of TWEAK. Bottom rows: microglia expressing 5 mRNAs of TWEAK. Scale bars, 5 μ m.

(C) Scatterplot of TWEAK expression (x axis) and proximal retinogeniculate synapses (y axis).

(D) Bar graph displaying the number of proximal retinogeniculate synapses (y axis) corresponding with increasing expression of TWEAK mRNA volume, binned as shown.

(legend continued on next page)

that STRING (search tool for the retrieval of interacting genes/proteins) analysis suggests make up a functional interaction network (Figure 5L). Most of these candidates are known to localize to synapses and play roles in synaptic function, particularly membrane trafficking, neurotransmitter receptor recycling, and cytoskeletal dynamics (Table S1). Altogether, these data suggest that TWEAK is recruited to retinogeniculate synapses by Fn14, where TWEAK signals locally through Fn14 to restrict the number of bulbous spines.

Synapses Near TWEAK-Expressing Microglia Are Preferentially Lost

What factors determine which synapses are removed through TWEAK-Fn14 signaling and which synapses are strengthened by Fn14 in the absence of TWEAK? Because TWEAK is a soluble factor, we reasoned that synapses proximal to microglia that express TWEAK may be preferentially targeted for TWEAK-dependent removal. To test this possibility, we developed an *in vitro* co-culture system in which neurons are dissociated from the embryonic WT mouse thalamus and co-cultured with microglia derived from TWEAK KO or WT littermate mice. These mice had been crossed previously to the Cx3cr1-GFP line, allowing us to visualize microglia in the TWEAK KO and WT co-cultures (Figures 6A, 6B, and S7; Jung et al., 2000). We found that addition of microglia to mCherry-labeled thalamic neurons leads to a significant decrease in the total number of spines that are present on neurons regardless of whether the microglia expressed TWEAK (Figure 6C). However, when we specifically looked at bulbous spines, we found that this class of spines was unaffected by addition of TWEAK KO microglia despite a 54% decrease in bulbous spines observed upon addition of WT microglia (Figures 6D–6F). This TWEAK-dependent decrease in bulbous spines depended on the proximity of the microglia to spines because we observed a significantly greater average distance between bulbous spines and the nearest TWEAK-expressing microglial cell compared with the distance between bulbous spines and non-TWEAK-expressing microglia (Figures 6G and 6H). In contrast, the distance between non-TWEAK-expressing microglia and thin spines was the same as for TWEAK-expressing microglia (Figures 6I and 6J). These data suggest that microglial release of TWEAK provides a local signal that restricts the number of bulbous spines on nearby thalamic relay neurons.

To complement our *in vitro* co-culture experiments, we next combined FISH with protein immunostaining to analyze the relationship between the amount of TWEAK mRNA expressed by a microglial cell and the proximity of retinal inputs to that same cell in dLGNs of visually stimulated mice. This approach takes advantage of the observation that microglia are heterogeneous in their expression of TWEAK so that some express high levels and some express low levels of TWEAK, even in the context of

visual stimulation (Figure 3D). Stimulated emission depletion (STED) microscopy indicated that microglia are commonly found within 50–200 nm of retinogeniculate synapses (Figure 7A); therefore, we designed our strategy to measure the number of retinal inputs within ~200 nm of a given microglia.

To determine whether sensory-induced microglial TWEAK regulates the number of nearby retinogeniculate synapses, we analyzed dLGN tissue from mice that were dark-reared and then acutely re-exposed to light to induce TWEAK expression in a subset of microglia. Remarkably, this analysis revealed a significant negative correlation between the level of TWEAK expressed by a microglial cell and the number of retinal synapses proximal to the microglial cell (Figures 7B and 7C; Pearson's coefficient $R = -0.4357$; $p < 0.001$). This correlation was also apparent when we plotted proximal inputs (y axis) versus microglial expression of TWEAK in bins (x axis) (Figures 7D and 7E) and when we divided microglia into low expressers and high expressers based on whether their TWEAK expression level fell below or above the median (Figures 7F and 7G). These data support the conclusion that, in response to visual stimulation, a subset of microglia express TWEAK, which, in turn, binds Fn14 on the surface of nearby thalamic neurons to locally suppress the number of retinogeniculate synapses.

DISCUSSION

Proposed Model of TWEAK-Fn14 Function in Synaptic Refinement

Based on our findings, we propose a mechanism where Fn14 differentially strengthens some synapses and eliminates others (Figure 8A). In this model, during the visual experience-dependent period of retinogeniculate circuit maturation, induction of Fn14 promotes formation and/or maintenance of bulbous spines on thalamic relay neurons, increasing the number of RGC-relay neuron connections and ultimately contributing to synapse strengthening within this neural circuit. However, when a retinogeniculate connection is proximal to TWEAK-expressing microglia, microglial TWEAK binds neuronal Fn14 to inhibit its ability to promote spine formation/maintenance, leading to spatially localized suppression of bulbous spine numbers. Thus, retinogeniculate synapses that are near TWEAK-expressing microglia remain weak and are ultimately eliminated specifically during the vision-sensitive phase of retinogeniculate circuit maturation. Taken together, these findings provide a potential explanation for how the protein product of an experience-induced gene such as Fn14 can effect strengthening of some synapses and elimination of others within the same neuron. They also define a role of microglia in synapse development that goes beyond phagocytic engulfment and is uniquely shaped by sensory experience (Figure 8B).

(E) Bar graph displaying the number of proximal retinogeniculate synapses (y axis) corresponding with increasing number of TWEAK mRNA transcripts, binned as shown.

(F) Comparison of proximal synapses in microglia expressing less than the median level of TWEAK (low TWEAK) versus those expressing greater than the median level of TWEAK (high TWEAK) assigned based on microglial volume occupied by mRNA signal.

(G) Same as (F) but mRNA plotted as transcripts per microglia. Values are normalized to microglial volume.

In (C), Pearson's correlation coefficient $R = -0.4357$. *** $p < 0.001$. In (F) and (G), Student's t test; * $p < 0.05$, *** $p < 0.001$; means are plotted with individual data points \pm SEM. In (D) and (E), average proximal input value across microglia per bin.




Figure 8. Model of TWEAK/Fn14-Dependent Synapse Regulation during Experience-Dependent Refinement

(A) Schematic of retinal inputs (orange) converging onto the dendrites of a relay neuron (teal). Alone, Fn14 increases bulbous spines to strengthen and maintain synapses, whereas TWEAK binding at other synapses leads to their ultimate disassembly. In the absence of experience, neither TWEAK nor Fn14 is expressed, so neither of these processes occur, and synapses remain in a weakened state but are not properly removed.

(B) We propose that the SD period of postsynaptic regulation by microglia identified in this study constitutes a later phase of microglia-driven circuit sculpting that follows earlier phases of phagocytic pruning and is driven by distinct molecular mechanisms.

SD Synapse Elimination Is Mechanistically Distinct from Phagocytic Engulfment

Since the discovery that dLGN relay neurons induce expression of MHC class I molecules in response to spontaneous activity, it has been appreciated that immune signaling molecules are expressed in the brain and play various roles in neural development (Corriveau et al., 1998). For example, microglia have been shown to engage the classical complement cascade to engulf less

active synapses during the first week of postnatal life, consistent with the idea that immune signaling pathways function in the brain at least in part in response to changes in neural activity (Gunner et al., 2019; Schafer et al., 2012). Notably, we found that the complement protein C1qa, which is required for activity-dependent engulfment at P5, does not regulate spines at P27 (Figure S8), indicating that this early phagocytosis-based process of microglial synapse removal is mechanistically distinct

from the SD regulation of synaptic structure we describe. It makes sense that, in the context of development, removal of the presynaptic input would precede disassembly of the postsynaptic site because performing these functions in reverse would potentially lead to excessive glutamate spillover, resulting in excitotoxicity. Our findings raise the intriguing question of whether the same microglial cell that engulfs presynaptic terminals might later remodel the postsynaptic specialization or whether different populations of microglia carry out these distinct processes. That TWEAK is induced in a subset of microglia suggests that different populations of microglia may be tuned to selectively engulf or remodel synapses. In any case, our results suggest that microglial roles in retinogeniculate synapse development extend beyond the phagocytic pruning of synapses that occurs prior to eye opening (Figure 8B).

Roles of Microglia in Regulating Synaptic Structure outside of the dLGN

Our findings regarding the function of microglia in the dLGN are complemented by recent studies of microglia in other brain regions. For example, in the hippocampus, microglia regulate spine structure without engulfing spines (Weinhard et al., 2018). Additionally, live imaging studies of microglial dynamics in the cortex have shown that microglia survey and contact synapses in response to sensory experience and that these microglia/synapse interactions influence the number, morphology, and physiology of spines (Akiyoshi et al., 2018; Tremblay et al., 2010). In a related study, Parkhurst et al. (2013) showed that depleting microglia from the mouse brain limits the spine turnover that usually occurs with motor learning. Most of these studies used live imaging of spines to study spine turnover and dynamics in the brains of living mice, an approach that will be useful for determining the mechanisms by which the TWEAK-Fn14 pathway regulates spines *in vivo*.

Implications for Human Disease

Brain disorders ranging from autism to Alzheimer's disease are characterized by changes in the numbers and structure of spines (Faludi and Mirnics, 2011; Glantz and Lewis, 2000; Hammond et al., 2019). These disorders are increasingly thought to involve the misregulation of microglial function (Hong et al., 2016; Velmeshchev et al., 2019). Our finding that microglia, via TWEAK-Fn14 signaling, shape the numbers and structure of spines during key periods of sensory experience-dependent circuit maturation raises the possibility that misregulation of this feature of microglial signaling may contribute to these disorders. Further examination of microglial roles in postsynaptic remodeling may therefore provide much-needed insight into how synapses become altered in pathological states that involve misregulation of microglial function.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY

- Lead Contact
- Materials Availability
- Data and Code Availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Generation of Fn14^{fl/fl}, Fn14^{fl/fl}; VGLUT2-Cre, and Fn14^{fl/fl}; Chx10-Cre mice
 - Generation of TWEAK^{fl/fl} and TWEAK^{fl/fl}; Cx3cr1-Cre mice
 - GridTape serial transmission electron microscopy
 - Golgi-staining and dendritic spine analysis
 - Protein detection by ELISA
 - Single-molecule fluorescence *in situ* hybridization (RNAscope)
 - Combined FISH and Immunofluorescence (FISH-IF)
 - Quantification of FISH-IF experiments
 - Synaptic engulfment and eye-specific segregation assays
 - Biochemical isolation of microglia-enriched fractions
 - Quantitative PCR
 - Microglial number and morphology analysis
 - Bilateral AAV injections into the dLGN
 - TWEAK localization experiment
 - Immunofluorescence
 - Fluorescence imaging
 - Synaptosomal fractionation
 - Immunoprecipitation
 - Mass spectrometry
 - *In vitro* thalamic neuron-microglia co-culture system
 - Statistical Analysis
 - Blinding
 - Statistical tests and software used

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.neuron.2020.08.002>.

ACKNOWLEDGMENTS

We thank Drs. C. Chen, E. Griffith, S. Ashrafi, E. Duffy Lacy, B. Kalish, and E. Pollina for thoughtful input on the manuscript. We thank Vance Soares for artistic work on the model. We thank Dr. Corey Harwell for Prkcd-Cre/Ai14 mouse tissue. We thank Dr. D.G.C. Hildebrand for help with sectioning and collection of dLGN tissue using GridTape. This work was supported by several core facilities at Harvard Medical School: the Neurobiology Imaging Facility, supported by NINDS P30 core center grant NS072030; the Vision Core, supported by grant P30 EY12196; the Taplin Mass Spectrometry Facility; the Genome Modification Facility; and the EM core at HMS. This work was supported by K99 MH120051 (NIMH) and a Goldenson postdoctoral fellowship (to L.C.), R01 NS028829 (NINDS) (to M.E.G.), NIH grants R21NS085320 and RF1MH114047 (to W.-C.A.L.), the Bertarelli Program in Translational Neuroscience and Neuroengineering, the Edward R. and Anne G. Lefler Center, and the Stanley and Theodora Feldberg Fund.

AUTHOR CONTRIBUTIONS

L.C. and M.E.G. conceptualized the study. L.C., B.S., L.C.B., W.-C.A.L., and M.E.G. designed experiments. L.C., S.A.R., J.S.P., and K.A.E. performed experiments. K.A.E. and L.C.B. provided Fn14 KO, TWEAK KO, and Fn14^{fl/fl} mice as well as the AAV-CASI-sTWEAK and AAV-CASI-mCherry viruses. All authors provided feedback on the manuscript, which was written by L.C. and M.E.G.

DECLARATION OF INTERESTS

L.C.B. and K.A.E. are employees and shareholders of Biogen.

Received: July 22, 2019

Revised: May 10, 2020

Accepted: August 5, 2020

Published: September 14, 2020

SUPPORTING CITATIONS

The following references appear in the Supplemental Information: Assali et al. (2017), Choi et al. (2006), Deng et al. (2011), Fischer von Mollard et al. (1994), Fukuda (2003), Gerges et al. (2005), Graf et al. (2009), Gromova et al. (2018), Han et al. (2009), Han et al. (2011), Kaeser et al. (2012), Kim et al. (2015), Park et al. (2018), Raemaekers et al. (2012), Sheehan and Waites (2019), Sheehan et al. (2016), Stahl et al. (1994), Teodoro et al. (2013), Wang et al. (2000), Wang et al. (2018), Zhao et al. (2014).

REFERENCES

- Akiyoshi, R., Wake, H., Kato, D., Horiuchi, H., Ono, R., Ikegami, A., Haruwaka, K., Omori, T., Tachibana, Y., Moorhouse, A.J., and Nabekura, J. (2018). Microglia Enhance Synapse Activity to Promote Local Network Synchronization. *eNeuro* 5, ENEURO.0088-18.2018.
- Assali, A., Le Magueresse, C., Bennis, M., Nicol, X., Gaspar, P., and Rebsam, A. (2017). RIM1/2 in retinal ganglion cells are required for the refinement of ipsilateral axons and eye-specific segregation. *Sci. Rep.* 7, 3236.
- Berry, K.P., and Nedivi, E. (2017). Spine Dynamics: Are They All the Same? *Neuron* 96, 43–55.
- Cardona, A.E., Huang, D., Sasse, M.E., and Ransohoff, R.M. (2006). Isolation of murine microglial cells for RNA analysis or flow cytometry. *Nat. Protoc.* 1, 1947–1951.
- Cheadle, L., Tzeng, C.P., Kalish, B.T., Harmin, D.A., Rivera, S., Ling, E., Nagy, M.A., Hrvatin, S., Hu, L., Stroud, H., et al. (2018). Visual Experience-Dependent Expression of Fn14 Is Required for Retinogeniculate Refinement. *Neuron* 99, 525–539.e0.
- Chen, C., and Regehr, W.G. (2000). Developmental remodeling of the retinogeniculate synapse. *Neuron* 28, 955–966.
- Chicheportiche, Y., Bourdon, P.R., Xu, H., Hsu, Y.M., Scott, H., Hession, C., Garcia, I., and Browning, J.L. (1997). TWEAK, a new secreted ligand in the tumor necrosis factor family that weakly induces apoptosis. *J. Biol. Chem.* 272, 32401–32410.
- Choi, S., Ko, J., Lee, J.R., Lee, H.W., Kim, K., Chung, H.S., Kim, H., and Kim, E. (2006). ARF6 and EFA6A regulate the development and maintenance of dendritic spines. *J. Neurosci.* 26, 4811–4819.
- Colonnier, M., and Guillery, R.W. (1964). Synaptic Organization in the Lateral Geniculate Nucleus of the Monkey. *Z. Zellforsch. Mikrosk. Anat.* 62, 333–355.
- Coriveau, R.A., Huh, G.S., and Shatz, C.J. (1998). Regulation of class I MHC gene expression in the developing and mature CNS by neural activity. *Neuron* 21, 505–520.
- Cowan, M., and Petri, W.A., Jr. (2018). Microglia: Immune Regulators of Neurodevelopment. *Front. Immunol.* 9, 2576.
- Datwani, A., McConnell, M.J., Kanold, P.O., Micheva, K.D., Busse, B., Shamloo, M., Smith, S.J., and Shatz, C.J. (2009). Classical MHC I molecules regulate retinogeniculate refinement and limit ocular dominance plasticity. *Neuron* 64, 463–470.
- Deng, L., Kaeser, P.S., Xu, W., and Südhof, T.C. (2011). RIM proteins activate vesicle priming by reversing autoinhibitory homodimerization of Munc13. *Neuron* 69, 317–331.
- Dohi, T., Borodovsky, A., Wu, P., Shearstone, J.R., Kawashima, R., Runkel, L., Rajman, L., Dong, X., Scott, M.L., Michaelson, J.S., et al. (2009). TWEAK/Fn14 pathway: a nonredundant role in intestinal damage in mice through a TWEAK/intestinal epithelial cell axis. *Gastroenterology* 136, 912–923.
- Faludi, G., and Mirics, K. (2011). Synaptic changes in the brain of subjects with schizophrenia. *Int. J. Dev. Neurosci.* 29, 305–309.
- Feinberg, I. (1982–1983). Schizophrenia: caused by a fault in programmed synaptic elimination during adolescence? *J. Psychiatr. Res.* 17, 319–334.
- Fischer von Mollard, G., Stahl, B., Li, C., Südhof, T.C., and Jahn, R. (1994). Rab proteins in regulated exocytosis. *Trends Biochem. Sci.* 19, 164–168.
- Fukuda, M. (2003). Distinct Rab binding specificity of Rim1, Rim2, rabphilin, and Noc2. Identification of a critical determinant of Rab3A/Rab27A recognition by Rim2. *J. Biol. Chem.* 278, 15373–15380.
- Gerges, N.Z., Brown, T.C., Correia, S.S., and Esteban, J.A. (2005). Analysis of Rab protein function in neurotransmitter receptor trafficking at hippocampal synapses. *Methods Enzymol.* 403, 153–166.
- Glantz, L.A., and Lewis, D.A. (2000). Decreased dendritic spine density on prefrontal cortical pyramidal neurons in schizophrenia. *Arch. Gen. Psychiatry* 57, 65–73.
- Graf, E.R., Daniels, R.W., Burgess, R.W., Schwarz, T.L., and DiAntonio, A. (2009). Rab3 dynamically controls protein composition at active zones. *Neuron* 64, 663–677.
- Graham, B.J., Hildebrand, D.G.C., Kuan, A.T., Maniates-Selvin, J.T., Thomas, L.A., Shanny, B.L., and Lee, W.-C.A. (2019). High-throughput transmission electron microscopy with automated serial sectioning. *bioRxiv* 657346, <https://doi.org/10.1101/657346>.
- Gromova, K.V., Muhia, M., Rothhammer, N., Gee, C.E., Thies, E., Schaefer, I., Kress, S., Kilimann, M.W., Shevchuk, O., Oertner, T.G., and Kneussel, M. (2018). Neurobeachin and the Kinesin KIF21B Are Critical for Endocytic Recycling of NMDA Receptors and Regulate Social Behavior. *Cell Rep.* 23, 2705–2717.
- Guillery, R.W., and Colonnier, M. (1970). Synaptic patterns in the dorsal lateral geniculate nucleus of the monkey. *Z. Zellforsch. Mikrosk. Anat.* 103, 90–108.
- Gunner, G., Cheadle, L., Johnson, K.M., Ayata, P., Badimon, A., Mondo, E., Nagy, M.A., Liu, L., Bemiller, S.M., Kim, K.W., et al. (2019). Sensory lesioning induces microglial synapse elimination via ADAM10 and fractalkine signaling. *Nat. Neurosci.* 22, 1075–1088.
- Hammond, T.R., Marsh, S.E., and Stevens, B. (2019). Immune Signaling in Neurodegeneration. *Immunity* 50, 955–974.
- Hamos, J.E., Van Horn, S.C., Raczkowski, D., and Sherman, S.M. (1987). Synaptic circuits involving an individual retinogeniculate axon in the cat. *J. Comp. Neurol.* 259, 165–192.
- Han, K., Kim, M.H., Seeburg, D., Seo, J., VerPELLI, C., Han, S., Chung, H.S., Ko, J., Lee, H.W., Kim, K., et al. (2009). Regulated RaiBP1 binding to RaiA and PSD-95 controls AMPA receptor endocytosis and LTD. *PLoS Biol.* 7, e1000187.
- Han, Y., Kaeser, P.S., Südhof, T.C., and Schneggenburger, R. (2011). RIM determines Ca²⁺ channel density and vesicle docking at the presynaptic active zone. *Neuron* 69, 304–316.
- Haubensak, W., Kunwar, P.S., Cai, H., Ciochi, S., Wall, N.R., Ponnusamy, R., Biag, J., Dong, H.W., Deisseroth, K., Callaway, E.M., et al. (2010). Genetic dissection of an amygdala microcircuit that gates conditioned fear. *Nature* 468, 270–276.
- Holtmaat, A., and Svoboda, K. (2009). Experience-dependent structural synaptic plasticity in the mammalian brain. *Nat. Rev. Neurosci.* 10, 647–658.
- Hong, Y.K., Park, S., Litvina, E.Y., Morales, J., Sanes, J.R., and Chen, C. (2014). Refinement of the retinogeniculate synapse by bouton clustering. *Neuron* 84, 332–339.
- Hong, S., Dissing-Olesen, L., and Stevens, B. (2016). New insights on the role of microglia in synaptic pruning in health and disease. *Curr. Opin. Neurobiol.* 36, 128–134.
- Hooks, B.M., and Chen, C. (2006). Distinct roles for spontaneous and visual activity in remodeling of the retinogeniculate synapse. *Neuron* 52, 281–291.
- Hooks, B.M., and Chen, C. (2008). Vision triggers an experience-dependent sensitive period at the retinogeniculate synapse. *J. Neurosci.* 28, 4807–4817.

- Hooks, B.M., and Chen, C. (2020). Circuitry Underlying Experience-Dependent Plasticity in the Mouse Visual System. *Neuron* *106*, 21–36.
- Jakubowski, A., Ambrose, C., Parr, M., Lincecum, J.M., Wang, M.Z., Zheng, T.S., Browning, B., Michaelson, J.S., Baetscher, M., Wang, B., et al. (2005). TWEAK induces liver progenitor cell proliferation. *J. Clin. Invest.* *115*, 2330–2340.
- Jung, S., Aliberti, J., Graemmel, P., Sunshine, M.J., Kreutzberg, G.W., Sher, A., and Littman, D.R. (2000). Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol. Cell. Biol.* *20*, 4106–4114.
- Kaesler, P.S., Deng, L., Fan, M., and Südhof, T.C. (2012). RIM genes differentially contribute to organizing presynaptic release sites. *Proc. Natl. Acad. Sci. USA* *109*, 11830–11835.
- Kalish, B.T., Cheadle, L., Hrvatin, S., Nagy, M.A., Rivera, S., Crow, M., Gillis, J., Kirchner, R., and Greenberg, M.E. (2018). Single-cell transcriptomics of the developing lateral geniculate nucleus reveals insights into circuit assembly and refinement. *Proc. Natl. Acad. Sci. USA* *115*, E1051–E1060.
- Katz, L.C., and Shatz, C.J. (1996). Synaptic activity and the construction of cortical circuits. *Science* *274*, 1133–1138.
- Kim, Y., Lee, S.E., Park, J., Kim, M., Lee, B., Hwang, D., and Chang, S. (2015). ADP-ribosylation factor 6 (ARF6) bidirectionally regulates dendritic spine formation depending on neuronal maturation and activity. *J. Biol. Chem.* *290*, 7323–7335.
- Maniates-Selvin, J.T., Grant, D., Hildebrand, D.G.C., Graham, B.J., Kuan, A.T., Thomas, L.A., Nguyen, T., Buhmann, J., Azevedo, A.W., et al. (2020). Reconstruction of motor control circuits in adult *Drosophila* using automated transmission electron microscopy. *bioRxiv*. <https://doi.org/10.1101/2020.01.10.902478>.
- Morgan, J.L., Berger, D.R., Wetzel, A.W., and Lichtman, J.W. (2016). The Fuzzy Logic of Network Connectivity in Mouse Visual Thalamus. *Cell* *165*, 192–206.
- Neniskyte, U., and Gross, C.T. (2017). Errant gardeners: glial-cell-dependent synaptic pruning and neurodevelopmental disorders. *Nat. Rev. Neurosci.* *18*, 658–670.
- Paolicelli, R.C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T.A., Guiducci, E., Dumas, L., et al. (2011). Synaptic pruning by microglia is necessary for normal brain development. *Science* *333*, 1456–1458.
- Parajuli, L.K., Tanaka, S., and Okabe, S. (2017). Insights into age-old questions of new dendritic spines: From form to function. *Brain Res. Bull.* *129*, 3–11.
- Park, D., Lee, U., Cho, E., Zhao, H., Kim, J.A., Lee, B.J., Regan, P., Ho, W.K., Cho, K., and Chang, S. (2018). Impairment of Release Site Clearance within the Active Zone by Reduced SCAMP5 Expression Causes Short-Term Depression of Synaptic Release. *Cell Rep.* *22*, 3339–3350.
- Parkhurst, C.N., Yang, G., Ninan, I., Savas, J.N., Yates, J.R., 3rd, Lafaille, J.J., Hempstead, B.L., Littman, D.R., and Gan, W.B. (2013). Microglia promote learning-dependent synapse formation through brain-derived neurotrophic factor. *Cell* *155*, 1596–1609.
- Peters, A., and Kaiserman-Abramof, I.R. (1970). The small pyramidal neuron of the rat cerebral cortex. The perikaryon, dendrites and spines. *Am. J. Anat.* *127*, 321–355.
- Raemaekers, T., Peric, A., Baatsen, P., Sannerud, R., Declerck, I., Baert, V., Michiels, C., and Annaert, W. (2012). ARF6-mediated endosomal transport of Telencephalin affects dendritic filopodia-to-spine maturation. *EMBO J.* *31*, 3252–3269.
- Rafols, J.A., and Valverde, F. (1973). The structure of the dorsal lateral geniculate nucleus in the mouse. A Golgi and electron microscopic study. *J. Comp. Neurol.* *150*, 303–332.
- Rowan, S., and Cepko, C.L. (2004). Genetic analysis of the homeodomain transcription factor Chx10 in the retina using a novel multifunctional BAC transgenic mouse reporter. *Dev. Biol.* *271*, 388–402.
- Saalfeld, S., Cardona, A., Hartenstein, V., and Tomancak, P. (2009). CATMAID: collaborative annotation toolkit for massive amounts of image data. *Bioinformatics* *25*, 1984–1986.
- Schafer, D.P., Lehrman, E.K., Kautzman, A.G., Koyama, R., Mardinly, A.R., Yamasaki, R., Ransohoff, R.M., Greenberg, M.E., Barres, B.A., and Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron* *74*, 691–705.
- Schneider-Mizell, C.M., Gerhard, S., Longair, M., Kazimiers, T., Li, F., Zwart, M.F., Champion, A., Midgley, F.M., Fetter, R.D., Saalfeld, S., and Cardona, A. (2016). Quantitative neuroanatomy for connectomics in *Drosophila*. *eLife* *5*, e12059.
- Sheehan, P., and Waites, C.L. (2019). Coordination of synaptic vesicle trafficking and turnover by the Rab35 signaling network. *Small GTPases* *10*, 54–63.
- Sheehan, P., Zhu, M., Beskow, A., Vollmer, C., and Waites, C.L. (2016). Activity-Dependent Degradation of Synaptic Vesicle Proteins Requires Rab35 and the ESCRT Pathway. *J. Neurosci.* *36*, 8668–8686.
- Sholl, D.A. (1953). Dendritic organization in the neurons of the visual and motor cortices of the cat. *J. Anat.* *87*, 387–406.
- Sorra, K.E., and Harris, K.M. (2000). Overview on the structure, composition, function, development, and plasticity of hippocampal dendritic spines. *Hippocampus* *10*, 501–511.
- Stahl, B., von Mollard, G.F., Walch-Solimena, C., and Jahn, R. (1994). GTP cleavage by the small GTP-binding protein Rab3A is associated with exocytosis of synaptic vesicles induced by alpha-latrotoxin. *J. Biol. Chem.* *269*, 24770–24776.
- Stevens, B., Allen, N.J., Vazquez, L.E., Howell, G.R., Christopherson, K.S., Nouri, N., Micheva, K.D., Mehalow, A.K., Huberman, A.D., Stafford, B., et al. (2007). The classical complement cascade mediates CNS synapse elimination. *Cell* *131*, 1164–1178.
- Szkarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva, N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* *47* (D1), D607–D613.
- Teodoro, R.O., Pekkumaz, G., Nasser, A., Higashi-Kovtun, M.E., Balakireva, M., McLachlan, I.G., Camonis, J., and Schwarz, T.L. (2013). Ral mediates activity-dependent growth of postsynaptic membranes via recruitment of the exocyst. *EMBO J.* *32*, 2039–2055.
- Tremblay, M.E., Lowery, R.L., and Majewska, A.K. (2010). Microglial interactions with synapses are modulated by visual experience. *PLoS Biol.* *8*, e1000527.
- Velmeshev, D., Schirmer, L., Jung, D., Haeussler, M., Perez, Y., Mayer, S., Bhaduri, A., Goyal, N., Rowitch, D.H., and Kriegstein, A.R. (2019). Single-cell genomics identifies cell type-specific molecular changes in autism. *Science* *364*, 685–689.
- Wang, Y., Sugita, S., and Südhof, T.C. (2000). The RIM/NIM family of neuronal C2 domain proteins. Interactions with Rab3 and a new class of Src homology 3 domain proteins. *J. Biol. Chem.* *275*, 20033–20044.
- Wang, J., Lv, X., Wu, Y., Xu, T., Jiao, M., Yang, R., Li, X., Chen, M., Yan, Y., Chen, C., et al. (2018). Postsynaptic RIM1 modulates synaptic function by facilitating membrane delivery of recycling NMDARs in hippocampal neurons. *Nat. Commun.* *9*, 2267.
- Weinhard, L., di Bartolomei, G., Bolasco, G., Machado, P., Schieber, N.L., Neniskyte, U., Exiga, M., Vadisiute, A., Raggioli, A., Schertel, A., et al. (2018). Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. *Nat. Commun.* *9*, 1228.
- Wiesel, T.N., and Hubel, D.H. (1963). Effects of Visual Deprivation on Morphology and Physiology of Cells in the Cats Lateral Geniculate Body. *J. Neurophysiol.* *26*, 978–993.
- Wiley, S.R., and Winkles, J.A. (2003). TWEAK, a member of the TNF superfamily, is a multifunctional cytokine that binds the TweakR/Fn14 receptor. *Cytokine Growth Factor Rev.* *14*, 241–249.

Yona, S., Kim, K.W., Wolf, Y., Mildner, A., Varol, D., Breker, M., Strauss-Ayali, D., Viukov, S., Guillems, M., Misharin, A., et al. (2013). Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity* *38*, 79–91.

Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., and Gerig, G. (2006). User-guided 3D active contour segmentation of anatomical

structures: significantly improved efficiency and reliability. *Neuroimage* *31*, 1116–1128.

Zhao, H., Kim, Y., Park, J., Park, D., Lee, S.E., Chang, I., and Chang, S. (2014). SCAMP5 plays a critical role in synaptic vesicle endocytosis during high neuronal activity. *J. Neurosci.* *34*, 10085–10095.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-Fn14	Cell Signaling Technology	Cat # 4403S; RRID:AB_10693941
Guinea Pig anti-VGLUT2	Millipore	Cat # AB2251; RRID:AB_2665454
Rabbit anti-Iba1	Wako	Cat # 019-19741; RRID:AB_839504
Rabbit anti-P2ry12	Sigma	Cat # HPA013796; RRID:AB_1854884
Goat anti-TWEAK	R&D Systems	Cat # AF1237; RRID:AB_2206219
Rabbit anti-GAPDH	Sigma-Aldrich	Cat # G9545; RRID:AB_796208
Mouse anti-PSD95	In-house	Cheadle et al., 2018
Chicken anti-MAP2	Lifespan Biosciences	Cat # LS-C61805; RRID:AB_1509808
Rabbit anti-GFAP	Abcam	Cat # ab4674; RRID:AB_304558
Goat anti-guinea pig AlexaFluor 647	Molecular Probes	Cat # A-21450; RRID:AB_141882
Goat anti-rabbit AlexaFluor 488	Thermo Fisher	Cat # A-11001; RRID:AB_2534069
Goat anti-mouse AlexaFluor 555	Thermo Fisher	Cat # A-21428; RRID:AB_2535849
Donkey anti-goat AlexaFluor 488	Thermo Fisher	Cat # A-11055; RRID:AB_2534102
Goat anti-guinea pig AlexaFluor 488	Molecular Probes	Cat # A-11073; RRID:AB_2534117
Goat anti-rabbit Complete IRdye 800 CW	Li-Cor biosciences	Cat # 827-08365; RRID:AB_10796098
Chemicals, Peptides, and Recombinant Proteins		
DAPI Fluoromount-G	Southern Biotech	Cat # 0100-20
NuPAGE LDS Sample Buffer (4X)	Novex	Ref # NP0007
Syn-PER synaptic protein extraction reagent	Thermo Fisher	Cat # 87793
Paraformaldehyde, 16%	Electron Microscopy Sciences	Cat # 15710
Protein A dynabeads	Life Technologies	Cat # 10002D
Triton X-100	Sigma-Aldrich	Cat # X100
Trizol	Life Technologies	Ref # 15596026
Cholera toxin-B conjugated to 488	Life Technologies	Cat # C34775
Cholera toxin-B conjugated to 555	Life Technologies	Cat # C34776
Lipofectamine 2000	Life Technologies	Cat # 11668500
Critical Commercial Assays		
FD Rapid Golgistain kit	FD NeuroTechnologies, Inc.	Cat #PK401A
PowerUp SYBR Green Master Mix	Life Technologies	Cat #A25743
RNAscope Multiplexed Fluorescence Detection kit	ACDBio	Cat #320850
RNeasy Micro Kit	QIAGEN	Cat #74004
RNAscope Multiplex Fluorescent Reagent Kit v2	ACDBio	Cat #323100
Mouse TWEAK DuoSet ELISA	R&D Systems	Cat #DY1237
BCA protein assay	Thermo Fisher	Cat # 23225
Silverquest Silver Staining kit	Thermo Fisher	Cat # LC6070
Western Lightning ECL Pro	Perkin-Elmer	Cat # NEL121001EA
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	The Jackson Laboratory	000664; RRID:IMSR_JAX:000664
Mouse: B6.Tnfrsf12a ^{tm1(KO)Biogen} (Fn14 KO)	Jakubowski et al., 2005	N/A
Mouse: B6.Tnfrsf12 ^{tm1(KO)Biogen} (TWEAK KO)	Dohi et al., 2009	N/A
Mouse: B6.Tnfrsf12a ^{(fl/fl)Biogen}	This paper	N/A
Mouse: B6.Tnfrsf12 ^{(fl/fl)Gree/J}	This paper	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mouse: B6.129P2(Cg)- <i>Cx3cr1</i> ^{tm1Litt} /J	The Jackson Laboratory	005582; RRID:IMSR_JAX:005582
Mouse: <i>Slc17a6</i> ^{tm2(cre)Low} /J	The Jackson Laboratory	028863; RRID:IMSR_JAX:016963
Mouse: Tg(Prkcd-glc-1/CFP,-Cre)	Haubensak et al., 2010	N/A
Mouse: B6.Cg-Gt(ROSA)26Sor ^{tm14(CAG-tdTomato)Hze} /J	The Jackson Laboratory	007914; RRID:IMSR_JAX:007914
Mouse: Tg(Chx10-EGFP/cre,-ALPP) ^{2Clc} /J	The Jackson Laboratory	005105; RRID:IMSR_JAX:005105
Mouse: B6J.B6N(Cg)- <i>Cx3cr1</i> ^{tm1.1(cre)Jung} /J	The Jackson Laboratory	025524; RRID:IMSR_JAX:025524
Mouse: B6.C1qa ^(KO)	Lab of Beth Stevens	N/A
Oligonucleotides		
qPCR primer: <i>Gapdh</i> (Forward): GGGTGTGAACACAG AGAAATA	Origene	Cat #: MP205604
qPCR primer: <i>Gapdh</i> (Reverse): CTGTGGTCATGAG CCCTTC	Origene	Cat #: MP205604
qPCR primer: <i>Tnfsf12</i> (TWEAK) (Forward): GCTGGGC AACGCTGTCT	Biogen	N/A
qPCR primer: <i>Tnfsf12</i> (TWEAK) (Reverse): GCGGTCC TCTGCTGTCA	Biogen	N/A
qPCR: <i>Cx3cr1</i> (Forward): GAGCATCACTGACATCTACCTCC	Origene	Cat #: MP202408
qPCR: <i>Cx3cr1</i> (Reverse): AGAAGGCAGTCGTGAGCTTGCA	Origene	Cat #: MP202408
qPCR: <i>P2ry12</i> (Forward): CATTGACCGCTACCTGAAGACC	Origene	Cat #: MP212229
qPCR: <i>P2ry12</i> (Reverse): GCCTCCTGTTGGTGAGAATCATG	Origene	Cat #: MP212229
Recombinant DNA		
AAV9-CASI-sTWEAK	Biogen	N/A
AAV9-CASI-mCherry	Biogen	N/A
pCAG-mCherry	In-house	N/A
Software and Algorithms		
ImageJ	NIH	https://fiji.sc/ or https://imagej.nih.gov/ij/
Prism	Graphpad	version 7.0b; RRID:SCR_002798
NeuroLucida	Microbrightfield	RRID:SCR_001775
Imaris	Bitplane	ImarisColoc
Metamorph	Molecular Devices	Version 1.0
Odyssey infrared imaging system	Li-Cor biosciences	Version 3.0
CATMAID	Saalfeld et al., 2009	https://catmaid.readthedocs.io/en/stable/
iTK-SNAP	Yushkevich et al., 2006	http://www.itksnap.org/pmwiki/pmwiki.php
AlignTK	N/A	http://mmbios.pitt.edu/installation
STRING	Szklarczyk et al., 2019	https://string-db.org/
Other		
FISH probe: <i>C1qa</i> , Channel 3	ACDBio	Cat # 441221-C3
FISH probe: <i>Cx3cr1</i> , Channel 2	ACDBio	Cat # 314221-C2
FISH probe: <i>Olig1</i> , Channel 3	ACDBio	Cat # 480651-C3
FISH probe: <i>Aldh1l1</i> , Channel 2	ACDBio	Cat # 405891-C2
FISH probe: <i>Cldn5</i> , Channel 3	ACDBio	Cat # 491611-C3
FISH probe: <i>Tnfrsf12a</i> (<i>Fn14</i>), Channel 3	ACDBio	Cat # 505311-C3
FISH probe: <i>P2ry12</i> , Channel 2	ACDBio	Cat # 317601-C2
FISH probe: <i>Vglut2</i> , Channel 2	ACDBio	Cat # 319171-C2
FISH probe: <i>Tnfsf12</i> (TWEAK), Channel 1	ACDBio	Cat # 552051
FISH probe: <i>Aldoc</i> , Channel 3	ACDBio	Cat # 429531-C3

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Michael E. Greenberg (meg@hms.harvard.edu).

Materials Availability

All unique resources generated in this study are available from the Lead Contact with a completed Materials Transfer Agreement. B6.Tnfrsf12a^{tm1(KO)Biogen} (Fn14 KO; [Jakubowski et al., 2005](#)), B6.Tnfrsf12a^{(fl/fl)Biogen} mice, and B6.Tnfrsf12^{tm1(KO)Biogen} (TWEAK KO; [Dohi et al., 2009](#)) are subject to restrictions imposed in an MTA by Biogen (Cambridge, MA).

Data and Code Availability

The published article includes all datasets generated or analyzed during the study.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All animal experiments were performed in compliance with protocols approved by the Institutional Animal Care and Use Committee (IACUC) at Harvard Medical School. The following mouse lines were used in the study: C57BL/6J (the Jackson Laboratory, JAX:000664); B6.129P2(Cg)-Cx3cr1^{tm1Litt}/J (Cx3cr1-GFP; the Jackson Laboratory, JAX: 005582); B6.Tnfrsf12a^{tm1(KO)Biogen} (Fn14 KO) ([Jakubowski et al., 2005](#)); B6.Tnfrsf12^{tm1(KO)Biogen} (TWEAK KO) ([Dohi et al., 2009](#)); a novel Fn14^{fl/fl} mouse line (B6.Tnfrsf12a^{(fl/fl)Biogen}; targeting strategy and validation shown in Figure S3); a novel TWEAK^{fl/fl} mouse line (B6.Tnfrsf12^{(fl/fl)Gree/J}; targeting strategy and validation shown in Figure S3); *Slc17a6*^{tm2(cre)Lowl/J} (VGLUT2-Cre; the Jackson Laboratory, JAX:028863); Tg(Prkcd-glc-1/CFP,-Cre) mice backcrossed to a C57Bl6/J background ([Haubensak et al., 2010](#)) then crossed to B6.Cg-Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze/J} (Ai14; the Jackson Laboratory, JAX:07914); B6J.B6N(Cg)-Cx3cr1^{tm1.1(cre)Jung/J} (Cx3cr1-Cre; the Jackson Laboratory, JAX: 025524), the Tg(Chx10-EGFP/cre,-ALPP)^{2Clc/J} line (Chx10-Cre; the Jackson Laboratory, JAX:005105) ([Rowan and Cepko, 2004](#)), and B6.C1qa^(KO) (lab of Beth Stevens). B6.Tnfrsf12a^{tm1(KO)Biogen}, B6.Tnfrsf12^{tm1(KO)Biogen}, and B6.Tnfrsf12a^{(fl/fl)Biogen} mice were provided by co-authors Dr. Linda Burkly and Katelin Ennis at Biogen. Most analyses were performed on mice at P27, though the developmental spine analysis also included P10, P20, and P90 ages, and some cohorts of mice were injected with AAVs at P12 to allow for spine analysis at P27. Thalamic neurons were cultured from pups at E15.5, and microglia were acutely isolated from mice between P15 and P20. Both males and females were included in all analyses. We did not note an effect of sex on any of the parameters measured in the study.

In most studies, mice were housed under standard conditions according to a 12-hour light/dark cycle (normally reared, NR). For the late-dark-rearing (LDR) paradigm, pups were bred in-house and housed with moms under standard light/dark conditions until P20, at which time they were weaned and moved into the dark, light-proof chamber of a ventilated cabinet. At P27, mice were moved into a separate, well-lit compartment of the cabinet for re-exposure to light for eight or twelve hours. Conversely, dark-reared, unstimulated control animals were euthanized by isoflurane and the brain removed in the dark by an investigator using night-vision goggles.

METHOD DETAILS

Generation of Fn14^{fl/fl}, Fn14^{fl/fl}; VGLUT2-Cre, and Fn14^{fl/fl}; Chx10-Cre mice

Fn14^{fl/fl} mice were generated by Taconic for Biogen. The strategy involved the insertion of two LoxP sites flanking the 2nd – 4th exons. To remove Fn14 from thalamic relay neurons, Fn14^{fl/fl} mice were bred in-house with VGLUT2-Cre mice (JAX:028863). To remove Fn14 from retinal ganglion cells, mice were bred in-house with Chx10-Cre mice (JAX:005105). Note that, although VGLUT2 expression has been reported in the retina, FISH revealed remaining Fn14 expression in the retina but not the dLGNs of Fn14^{fl/fl}; VGLUT2-Cre mice. We therefore considered this mouse to be a dLGN-specific knockout.

Generation of TWEAK^{fl/fl} and TWEAK^{fl/fl}; Cx3cr1-Cre mice

TWEAK^{fl/fl} mice were designed to excise the 3rd coding exon by floxing this region with LoxP sites in the introns directly before and after exon three. This strategy was predicted to introduce a STOP codon and lead to nonsense-mediated decay of the message. LoxP sites were introduced by CRISPR technology at the Genome Modification Facility in Cambridge, Massachusetts. Microglia-specific loss of TWEAK in TWEAK^{fl/fl} mice bred to the microglial Cre driver Cx3cr1-Cre was validated by RNAscope. In addition, significant decreases (but not complete ablation) of TWEAK at the whole-tissue level was demonstrated by qPCR.

GridTape serial transmission electron microscopy

Tissue preparation

An Fn14 KO mouse and a WT littermate at P27 were *trans*-cardially perfused with 2% paraformaldehyde and 2.5% glutaraldehyde in 0.1 M Cacodylate buffer with 0.04% CaCl₂. Coronal sections of the brain containing the dLGN (300 μm thickness) were cut on a Vibratome (VT1000S, Leica) and stained with 1% osmium tetroxide and 1.5% potassium ferrocyanide followed by 1% uranyl acetate,

then lead aspartate. Sections were dehydrated with a graded ethanol series and embedded in epoxy resin (TAAB 812 Epon, Cane MCO).

Sectioning and collection

For both the WT and Fn14 KO samples, 250 ultrathin sections (~45 nm) of the dLGN tissue were cut using an ultramicrotome (UC7, Leica) with a diamond knife (Diatome). Sections were collected onto GridTape using a modified automated tape-collecting ultramicrotome (ATUM) equipped with an optical interrupter (GP1A57HRJ00F, Sharp) and a hall-effect sensor (A1301EUA-T, Allegro MicroSystems) and magnet. The tape's movement speed was varied based on signals from the interrupter and hall-effect sensor to consistently place sections of the dLGN over GridTape's slots (Graham et al., 2019).

Post-section staining and TEM imaging

Following sectioning and collection, sample contrast was enhanced via post-section staining with Reynolds lead citrate (Ultrastain II, Leica). Electron microscopy imaging of the collected thin sections from both samples was performed at 4.3 nm pixel size on a custom JEOL 1200EX transmission electron microscope modified to hold and automatically image reels of GridTape (Graham et al., 2019). After acquisition, images were stitched and aligned with AlignTK (<http://mmbios.pitt.edu/installation>) into three-dimensional (3D) volumes.

Quantification of bouton properties

Aligned volumes were uploaded to CATMAID (Saalfeld et al., 2009; Schneider-Mizell et al., 2016) for manual annotation of retinal boutons, their synapses, and the type of the postsynaptic structure (i.e., shaft, bulbous spine, thin spine, or broad non-bulbous spine). The dLGNs were divided into 4 quadrants to ensure synapses were analyzed across all regions for both Fn14 KO and WT sections. The proportions of retinal inputs converging onto spines versus dendritic shafts was determined by randomly sampling 250 boutons per quadrant (1,000 total) for each condition. Retinal boutons were identified based upon morphological properties (Colonnier and Guillery, 1964; Guillery and Colonnier, 1970). Specifically, retinal boutons were identified by their mitochondria, which are uniquely large and pale, in combination with the relatively large size of the boutons themselves. 3D reconstructions of selected boutons (as seen in Figures 1C and 1D) were performed in ITK-SNAP (<http://www.itksnap.org/pmwiki/pmwiki.php>; Yushkevich et al., 2006) and Gaussian image smoothing was applied according to the following parameters: standard deviation, 0.90; approximation error max, 0.10. All analyses were performed with the researcher blinded to condition.

Golgi-staining and dendritic spine analysis

Golgi staining was performed with the FD Rapid GolgiStain kit (FC Neurotechnologies, Inc) according to the manufacturer's protocol. Following sample processing, dendritic segments were traced in x, y, and z planes using a Zeiss Axioskop microscope (63X objective) and Neurolucida (Microbrightfield Bioscience). To categorize spines by morphology, we defined bulbous spines as those whose head diameter is at least 2X that of its neck, spines with a relatively uniform width (head to neck width ratio less than 2) of less than 0.5 μm as thin spines, and spines with a relatively uniform width of greater than 0.5 μm as broad non-bulbous spines. In a majority of cases, head-to-neck ratio cutoffs were clear by eye after some analysis on training datasets by the investigator. In cases where this was not clear, the criteria were determined by measurement. We find by hierarchical clustering that these morphological subtypes may represent distinct biological categories (Figures S2B and S2C). In addition to these categorizations, we also measured the spine head diameter and the length of spines overall regardless of classification. Although spines are thought to exist along a structural continuum, complementing our measurements of spine structure overall with an analysis of spine subtypes is useful in allowing our analysis to be compared with analyses in other brain regions, many of which describe spine morphology according to subclasses similar to those used in this study (Peters and Kaiserman-Abraham, 1970; Sorra and Harris, 2000). Example images of Golgi-stained spines were obtained on an Olympus BX63 microscope with a 100X objective.

Protein detection by ELISA

The dLGNs and primary visual cortices of normally reared, dark-reared, and visually stimulated mice were microdissected in ice-cold PBS following sectioning on a vibratome (VT1000S, Leica) and flash-frozen in liquid nitrogen. On the day of the assay, the tissues were thawed and resuspended in 500 μL of ice-cold RIPA buffer (Life Technologies) including complete protease inhibitor cocktail tablet (Roche) and phosphatase cocktails two and three (Sigma). Homogenization was performed by douncing 25X in a 2 mL douncer and the protein concentration of each sample was determined by BCA assay (Pierce). The samples were then diluted to achieve a concentration of 100 $\mu\text{g}/\mu\text{L}$ for each sample. TWEAK protein expression was measured by ELISA (R&D Systems, DY1237) according to the manufacturer's instructions, with 30 μg of sample loaded per well in triplicate.

Single-molecule fluorescence *in situ* hybridization (RNAscope)

The brains of euthanized mice were harvested and immediately embedded in OCT (Optimum Cutting Temperature) on dry ice then stored at -80°C . Twenty μm sections were cut on a Leica CM 1950 Cryostat, collected on Superfrost Plus slides, and stored at -80°C until the day of the experiment. smFISH was performed using RNAscope (Advanced Cell Diagnostics) according to the manufacturer's protocol for fresh-frozen sections and as described (Cheadle et al., 2018). Sections were probed for the microglial markers *Cx3cr1* (314221-C2), *C1qa* (441221-C3), and *P2ry12* (317601-C2). In addition, the excitatory relay neuron marker *VGLUT2* (319171-C2), the interneuron marker *Gad1* (400951-C2), the endothelial cell marker *Cldn5* (491611-C3), the

astrocyte markers *Aldh111* (405891-C2) and *Aldoc* (429531-C3), the oligodendrocyte marker *Olig1* (480651-C3), *TWEAK* (552051), and *Fn14* (505311-C3) were also used.

Blinded multi-channel images were imported into ImageJ. Following background subtraction, cell types were identified based on robust expression of at least eight marker gene mRNAs (*Cx3cr1* or *P2ry12* for microglia, *Cldn5* for endothelial cells, *Aldh111* or *Aldoc* for astrocytes, *VGLUT2* for relay neurons, *Gad1* for interneurons, and *Olig1* for oligodendrocytes), and the DAPI-stained nuclei of these cells were traced. Next, the channel representing *TWEAK* or *Fn14* signal was opened, and the number of *TWEAK* mRNAs per microglial cell body or *Fn14* per relay neuron was quantified.

Combined FISH and Immunofluorescence (FISH-IF)

We combined RNAscope FISH and immunofluorescence in order to visualize and measure the relationship between microglial *TWEAK* expression and retinogeniculate synapse number. Mice were perfused in 4% paraformaldehyde and post-fixed overnight at 4°C. Brains were cryoprotected by incubation in 15% then 30% sucrose. Sections of 20 μm were made and stored at –80°C until use.

Tissue was processed with the RNAscope Multiplex Fluorescent V2 Assay (Advanced Cell Diagnostics, #323100) according to the manufacturer's instructions. Several modifications to the manufacturer's protocol were made to optimize co-visualization of mRNA and protein signal. First, the pretreatment fixation in 4% PFA was performed for 20 minutes. Next, the hydrogen peroxide incubation step was 10 minutes. For the target retrieval step, the ACD target retrieval reagent was brought to a boil and allowed to cool to 90°C on the bench top, at which point sections were added and incubated for 15 minutes. Sections were incubated with Protease III for 20 minutes.

After performing all *TWEAK* mRNA probe detection steps, sections were blocked for 30 minutes at room temperature in PBS adjusted to 0.3% Triton X-100 and 5% normal goat serum. Sections were incubated at 4°C overnight with primary antibodies: rabbit anti-Iba1 (Wako, 019-19741; 1:500), rabbit anti-P2ry12 (Sigma, HPA013796; 1:500), and guinea pig anti-VGLUT2 (AB2251, Millipore; 1:500) in PBS adjusted to 0.1% Triton X-100 and 5% normal goat serum. Sections were washed in RNAscope wash buffer 2 times for 2 minutes per wash. Secondary Alexa Fluors (Invitrogen; 1:500) were used to detect primary antibodies via incubation with sections for 2 hours at RT. Sections were then washed and coverslipped in Fluoromount G + DAPI (ThermoFisher).

Quantification of FISH-IF experiments

Sections were imaged in three dimensions on a laser-scanning confocal microscope (Zeiss FV1000) with a 60X objective. After pre-processing to enhance contrast and subtract background in ImageJ, Z stacks of individual microglia containing all three channels were imported into Imaris (Bitplane). Using the "Surface" function, volumetric reconstructions were generated of (1) the microglia; (2) *TWEAK* mRNA signal; (3) retinogeniculate synapses (VGLUT2 signal); (4) "masked" *TWEAK* mRNA puncta within the boundaries of the microglial volume; and (5) "masked" retinogeniculate synapses co-localizing with the microglial volume. Since the resolution of confocal images is roughly 200 nm, we denoted colocalized synapses as those most proximal to the microglia in question. The amount of *TWEAK* within the microglial cell was measured based upon (1) the percentage of the microglial volume occupied by *TWEAK* signal, and (2) the number of discrete *TWEAK* puncta detected per microglial volume. Because in several cases individual *TWEAK* mRNA were so close together that they appeared as one continuous unit, we believe the volumetric measurement is most accurate. In our final quantifications, we not only normalized the amount of *TWEAK* mRNA detected to the volume of the microglia analyzed, we also normalized the amount of colocalizing retinogeniculate inputs to the microglial volume as well. Furthermore, because retinal synapse density varies across the dLGN, we normalized all inputs associated with the microglia from the same section to the overall average input density across the image.

Synaptic engulfment and eye-specific segregation assays

Retinal inputs to the dLGN were labeled through ocular injections of cholera toxin-B anterograde tracers conjugated to Alexa Fluor dyes (Invitrogen). Conversely, *Prkcd-Cre/LSL-TdTomato* animals in which relay neurons are labeled with TdTomato were analyzed. After 48 hours, mice were euthanized and perfused with 4% paraformaldehyde (PFA) then the brains were sectioned and immunostained for microglia using rabbit anti-Iba1 (Wako, 019-19741) and rabbit anti-P2ry12 (Sigma, HPA013796) in combination. Sections were imaged in three dimensions on a laser-scanning confocal microscope (Zeiss FV1000) with a 60X objective. Z stacks containing all three channels were imported into Imaris (Bitplane) and the volumes of individual microglia were reconstructed based upon Iba1 and P2ry12 staining. The volume of each microglial cell occupied by phagocytosed retinal inputs was quantified as described (Schafer et al., 2012).

For eye-specific segregation analysis, z stacks were imaged at 10X to allow visualization of the ipsi- and contralateral input regions across a coronal section of the dLGN. Images were imported into Metamorph (Molecular Devices). The fluorescence intensity of the labeled ipsilateral retinal inputs was increased to include all signal above background. Next, the intensity of the contralateral input signal was gradually increased from 0 to 250, and the overlap between the signals at each contralateral threshold was measured.

Biochemical isolation of microglia-enriched fractions

Microglia were purified from the visual cortices of dark-reared, unstimulated mice and mice re-exposed to light for eight or twelve hours as previously described (Cardona et al., 2006). After anesthetization with isoflurane, mice were perfused with ice cold 1X HBSS (GIBCO) and their brains dissected and placed in RPMI buffer (Sigma). After discarding the cerebellum, the brains were minced with a razor blade and transferred to a 2 mL dounce homogenizer containing 1.5 mL of RPMI buffer. The brains of eight mice per condition were pooled at this point and dounced 30X with the loose pestle then 25X with the tight pestle. Homogenates were transferred to a 15 mL falcon tube containing 5.5 mLs of RPMI (total volume, 7 mL). Standard isotonic percoll (SIP; 4.5 mL percoll with 0.5 mL 10X HBSS) was then added to the homogenate and mixed by pipetting. The homogenate was then overlaid onto 3 mLs of 70% SIP in a 15 mL falcon tube and centrifuged at 500 \times g for 30 minutes, 18°C. After removal of the lipid layer at the top, the microglial fraction was recovered from the 30%/70% SIP interface (about 1 mL) and transferred to a falcon tube containing 10 mL HBSS. Following centrifugation at 900 \times g for 7 minutes at 18°C, the supernatant was discarded and pelleted microglia were resuspended in Trizol (Sigma), flash-frozen in liquid nitrogen, and stored at -80°C .

Quantitative PCR

Microglia-enriched fractions from visual cortex flash-frozen in Trizol were thawed and purified using the RNeasy Micro kit with on-column DNase digestion (QIAGEN). Reverse transcription was performed using the Superscript III first-strand synthesis kit (Invitrogen). Real-time quantitative PCR was performed using SYBR green mix (Life Technologies). Levels of endogenous *TWEAK* were detected with the following primers: GCTGGGCAACGCTGTCT (F) and GCGGTCCTCTGCTGTCA (R), and normalized to the expression of the microglial marker *Cx3cr1*: GAGCATCACTGACATCTACCTCC (F) and AGAAGGCAGTCGTGAGCTTGCA (R). Validation of loss of *TWEAK* in *TWEAK^{fl/fl}*; *Cx3cr1-Cre* was performed by the same qPCR approach on whole-brain tissue with *TWEAK* levels normalized to *GAPDH* and *P2ry12* expression using the following primers: *GAPDH* forward (GGGTGTGAACCACGAGAAATA), *GAPDH* reverse (CTGTGGTCATGAGCCCTTC), *P2ry12* forward (CATTGACCGCTACCTGAAGACC), and *P2ry12* reverse (GCCTCTGT TGGTGAGAATCATG).

Microglial number and morphology analysis

Microglia were visualized by immunostaining for Iba1 and P2ry12 and z stacks were taken on a laser-scanning confocal microscope (Zeiss FV1000). Images were imported into ImageJ, background subtracted, and the numbers of Iba1-positive cells were measured per unit dLGN area. For Sholl analysis, z stacks of individual microglia were opened in Neurolucida (Microbrightfield) and the cell bodies and primary protrusions were traced in 3D using a 40X objective. The tracings were then imported into Neurolucida Explorer and a Sholl analysis was performed to measure the morphological complexity of each microglial cell by quantifying the number of intersections between microglial processes and a series of concentric circles radiating from the soma outward (Sholl, 1953).

Bilateral AAV injections into the dLGN

AAV9-CAS1-sTWEAK and AAV9-CAS1-mCherry were designed, produced, and characterized by co-authors Dr. Linda Burkly and Katelyn Ennis at Biogen. We further confirmed the efficacy of the TWEAK-expressing virus using RNAscope as described above (see inset of Figure 4H). For injections, P12 mice were anesthetized with isoflurane and stereotaxically injected with 1 μL of AAV-sTWEAK (right dLGN) or AAV-mCherry (left dLGN) at the following stereotaxic coordinates: $-2.0y, +/ - 2.1x, 2.95z$ (mm). Animals were administered Flunixin twice daily for three days following the injections. Most mice appeared normal and comfortable within one day following injection. Injected mice were usually housed according to standard light/dark conditions until P27 at which point the animals were euthanized and the brains subjected to Golgi-staining. In some cases, mice were reared according to standard conditions until P20 and then late-dark-reared (LDR) until P27.

TWEAK localization experiment

TWEAK KO mice, *Fn14^{fl/fl}* mice, and *Fn14^{fl/fl}*; *VGLUT2-Cre* mice at P27 were perfused with ice cold oxygenated artificial cerebrospinal fluid (ACSF) containing (in mM): 125 NaCl, 26 NaHCO_3 , 1.25 NaH_2PO_4 , 2.5 KCl, 1.0 MgCl_2 , 2.0 CaCl_2 , and 25 glucose (Sigma), adjusted to 310-312 mOsm with water. Coronal sections of dLGN at 150 μm were sliced on a vibratome (VT1000S, Leica) and kept on ice in ACSF. Sections were incubated with either ACSF or ACSF containing 200 ng/mL of recombinant mouse TWEAK (Biogen) and rotated at 4°C for 2 hours. ACSF was aspirated and 500 μL of 4% PFA was added to fix slices for 20 minutes at room temperature. Sections were then washed in 1X PBS (GIBCO) 3x for 10 minutes per wash and blocked in PBS adjusted to 1.25% Triton X-100 (United States Biological) and 10% fetal bovine serum (GIBCO; IHC buffer) for 1 hour at room temperature. Sections were then stained in IHC buffer overnight at 4°C with guinea pig anti-VGLUT2 (AB2251, Millipore) and goat anti-TWEAK (AF1237, R&D Systems) at 1:1000. The next day, sections were washed 3 times in PBS then incubated in secondary Alexafluor Dyes (Molecular Probes) in IHC buffer at 1:500 for 1 hour. Sections were washed in PBS and mounted onto slides in Fluoromount G plus DAPI (SouthernBiotech). Imaging parameters given below.

Immunofluorescence

Fifty μm sections of 4% PFA-fixed brains were made on a vibratome and blocked in PBS adjusted to 1.25% Triton X-100 and 10% fetal bovine serum for one hour at room temperature. Primary antibody incubations were performed at 4°C overnight in blocking

buffer. The next day, sections were washed three times in PBS then probed with secondary Alexafluor dyes (Molecular Probes or Thermo Fisher) for one hour at room temperature. Sections were again washed three times in PBS and mounted onto Superfrost Plus slides (Fisher) using Fluoromount G plus DAPI (SouthernBiotech). Primary antibodies used include guinea pig anti-VGLUT2 (AB2251, Millipore), rabbit anti-Iba1 (Wako, 019-19741), rabbit anti-P2ry12 (Sigma, HPA013796), and rabbit anti-GFAP (Abcam, ab4674).

Fluorescence imaging

Imaging of immunofluorescence and FISH was performed on an Olympus FluoView 1000 laser scanning confocal microscope equipped with 405 nm, 440 nm, 488 nm, 515 nm, 559 nm, and 635 nm excitation lasers, and 10x air 0.4NA, 20x air 0.75NA, 40x oil 1.3NA, 60x oil 1.42NA, and 100x oil 1.4NA objectives. STED imaging was performed on a Leica Sp8 STED/FLIM system using a 100X oil objective and a white light laser providing continuous excitation tuning from 470 nm to 670 nm.

Synaptosomal fractionation

Synaptosomal fractionation was performed using Syn-PER synaptic protein extraction reagent (ThermoFisher, 87793) according to the manufacturer's instructions. Briefly, Complete protease inhibitor tablet (Roche) and phosphatase inhibitor cocktails 2 and 3 (Sigma) were added to 10 mL of ice cold Syn-PER reagent. The forebrains of 4 WT adult mice were minced and added to 2 mL of Syn-PER reagent in a 2 mL dounce homogenizer and homogenized ~10 times. The sample was then centrifuged at 1200xg for 10 minutes at 4°C. The supernatant was then centrifuged again at 15,000xg for 20 minutes at 4°C. The supernatant (cytosolic fraction) was then transferred to a separate tube and the synaptosomal pellet was resuspended in 200 μ L of Syn-PER reagent. The protein concentrations of homogenate, cytosolic non-synaptic fraction, and synaptosomal fraction were determined by BCA protein assay (Pierce) and 30 μ g of each were combined with protein sample buffer and run on a western blot as described below.

Immunoprecipitation

Validation of Fn14^{fl/fl} mice was performed by immunoprecipitating Fn14 from the whole forebrain using a rabbit anti-Fn14 antibody (Cell Signaling Technologies, 4403s) at 1:50 as described (Cheadle et al., 2018). Briefly, the brains of WT, Fn14^{fl/fl} mice, and Fn14^{fl/fl}; VGLUT2-Cre mice were homogenized in buffer containing 10 mM HEPES-KOH pH 7.5, 25 mM KAc, 320 mM sucrose, 1% Triton X-100, and 250 mM NaCl, along with a complete protease inhibitor cocktail tablet (Roche) and phosphatase cocktails two and three (Sigma). Following homogenization in a 2 mL douncer, homogenates were rotated at 4°C for 10 minutes and then centrifuged at 14,000 RPM for 15 minutes at 4°C. Supernatants were cleared by rotating for 1 hour with 50 μ L of protein A dynabeads (Invitrogen). Beads were collected on a magnet and supernatants transferred to a new eppendorf tube. Rabbit anti-Fn14 (4403s) was added to each sample at 1:50 and samples were rotated for 2 hours at 4°C. Samples were then rotated with 50 μ L of protein A dynabeads at 4°C which were then captured on a magnet. The supernatant was discarded and the beads were washed in 1 mL of wash buffer (homogenization buffer above but without sucrose) 4 times for 10 minutes/wash at 4°C. Proteins were then eluted from beads in 100 μ L 1X Nupage LDS 4X Sample Buffer (Life Technologies) with 10% 2-mercaptoethanol by boiling at 95°C for one minute.

For western blotting, samples were run on 12% Bis-tris gels (Life Technologies) and transferred to 0.2 μ m pore nitrocellulose (Bio-RAD). Blocking and antibody incubations were performed in TBS-T with 5% dry milk. Primary antibodies used were rabbit anti-Fn14 (4403s), guinea pig anti-VGLUT2 (AB2251, Millipore), mouse anti-PSD-95 (in-house), and rabbit anti-GAPDH (G9545, Sigma). Secondary antibodies included anti-rabbit, anti-guinea pig, and anti-mouse conjugated to HRP. Blots were developed with Western Lightning ECL Pro (Perkin-Elmer). In some cases, western blots were quantified using the Li-Cor Odyssey system and fluorescent secondaries (Li-Cor biosciences).

Mass spectrometry

For unbiased discovery of Fn14 interactors, Fn14 was immunoprecipitated from the brains of an adult WT mouse and an Fn14 KO littermate as described above. The immunoprecipitate was run on a bis-tris gel which was then silver stained using the SilverQuest kit from Invitrogen according to manufacturer's instructions. The protein bands were excised and submitted to the Taplin Biological Mass Spectrometry Facility where the samples were analyzed by LC-MS/MS. Briefly, gel pieces were trypsinized and dehydrated, then rehydrated in 50 mM ammonium bicarbonate solution containing 12.5 ng/ μ L modified sequencing-grade trypsin (Promega, Madison, WI) at 4°C. After 45 min, the excess trypsin solution was removed and replaced with 50 mM ammonium bicarbonate solution to just cover the gel pieces. Samples were then incubated at 37°C overnight and extracted by removing the ammonium bicarbonate solution, followed by one wash with a solution containing 50% acetonitrile and 1% formic acid. The extracts were then dried in a speed-vac for 1 hour.

Samples were reconstituted in 5 - 10 μ L of HPLC solvent A (2.5% acetonitrile, 0.1% formic acid). Each sample was loaded onto a Famos auto sampler (LC Packings, San Francisco, CA). Peptides were eluted with increasing concentrations of solvent B (97.5% acetonitrile, 0.1% formic acid). Eluted peptides were subjected to electrospray ionization and then entered into an LTQ Orbitrap Velos Pro ion-trap mass spectrometer (Thermo Fisher Scientific, Waltham, MA). Peptides were detected, isolated, and fragmented to produce a tandem mass spectrum of specific fragment ions for each peptide. Peptide sequences (and hence protein identity) were determined by matching protein databases with the acquired fragmentation pattern by the software program, Sequest (Thermo Fisher Scientific, Waltham, MA).

We performed 3 individual bioreplicates of the experiment. All peptide hits listed in the manuscript were identified in at least 2 of 3 bioreplicates and were never detected in IgG controls from the same wild-type mouse brains or in separate Fn14 co-IPs from the brains of Fn14 KO mice; $n = 3$ experiments from 3 mice per genotype. STRING analysis was performed to assess functional interactions between molecules identified as possible Fn14 interactors (STRING; <https://string-db.org/>; Szklarczyk et al., 2019).

In vitro thalamic neuron-microglia co-culture system

The night before dissections, glass coverslips were coated in Poly-D-Lysine (Sigma) and Laminin (Life Technologies) diluted in sterile, autoclaved water, and incubated overnight in a sterile cell culture incubator set to 37°C, 5% CO₂. The next morning, the coverslips were washed with sterile water 2 times and once with Neurobasal plain (GIBCO).

Prior to dissection, Dissociation Media (DM) was prepared as a 10X stock. 100 mM MgCl₂, 100 mM HEPES, and 10 mM kynurenic acid were added to HBSS w/o Mg²⁺ or Ca²⁺ (GIBCO). The solution was pH-ed to 7.2. 10X stocks were stored up to 6 months at -20°C, and 1X dilutions in HBSS were stored up to 1 month at 4°C.

In a tissue culture hood, 0.06 g of ovomucoid trypsin inhibitor (Worthington Biochemical) and 0.06 g of BSA (Sigma) were added to 6 mL of 1X DM in a falcon tube, which was labeled “Heavy.” The Heavy solution was warmed in a water bath at 37°C until both the inhibitor and BSA were dissolved (about 10 minutes). Heavy was pH-ed with sterile, filtered 1N NaOH from a P20 pipette tip to about 7.2 (~20 μL of 1N NaOH accomplished this). Next, 1 mL of this solution was added to 9 mL of 1X DM to make a solution labeled “Light.”

Pregnant female mice at E15.5 gestation day were euthanized by isoflurane and fetuses surgically removed from the amniotic sac. The fetuses were decapitated and the brains removed. The meninges were removed using fine forceps and the brains were bisected down the longitudinal fissure. Subcortical thalamic tissue was removed from each hemisphere and placed in 1X DM on ice until all fetuses had been processed. Two to three litters were typically pooled for each experiment.

Papain solution was prepared by combining 5 mL of 1X DM, a few grains of L-Cysteine (Sigma), and 100 μL of papain (10108014001, Sigma) and warming the solution in a water bath until dissolved. Next, the solution was filtered through a 0.02 μm filter into a separate falcon tube containing 2.5 mg of DNase (Sigma). The 1X DM in which the thalami were pooled was aspirated from the tube, and 3 mL of the papain/DNase solution was added to the tissue. The falcon tube was placed in the 37°C incubator for 10 minutes and stirred about every 2.5 minutes. The papain solution was aspirated and the tissue washed in 3 mL of Light solution 3 times, then 3 mL of Heavy solution 1 time. Next, the tissue was washed 1 time in 3 mL of Neurobasal complete [Neurobasal (GIBCO) adjusted to 1% Penicillin/Streptomycin, 1% Glutamax, and 10 mL B27 Supplement (1X)].

Tissue was dissociated by adding 2 mL of Neurobasal complete and pipetting 1 time with a P1000. Tissue chunks were allowed to settle then 1 mL of the supernatant was added to a clean falcon tube through a 50 μm cell strainer. One mL of fresh Neurobasal complete was added back to the tube with the tissue and this trituration/filtering step was repeated 10 to 15 times. Finally, the remaining solution was strained through the cell strainer and 3 mL of Neurobasal complete was added to the strainer to wash it. Cells were counted on a hemocytometer and plated at a concentration of 100,000 cells/well in 1 mL of Neurobasal complete. After incubating for about 1 hour, the media was changed to fresh pre-warmed Neurobasal complete. Media was supplemented about every 3 days by removing 300 μL of the media from each well and adding back 400 μL fresh Neurobasal. At 5 DIV, neurons were sparsely transfected with 1 μg of CAG-mCherry using Lipofectamine 2000 (ThermoFisher) according to the manufacturer’s instructions.

For neuron-microglia co-cultures, microglia were acutely isolated from the brains of Cx3cr1-GFP; TWEAK KO or Cx3cr1-GFP; TWEAK WT mice between P15 and P20 as described in the “Biochemical isolation of microglia-enriched fractions” section above, with the exception that, following the final centrifugation to pellet the microglia, the pellet was re-suspended in Neurobasal complete and the cells were counted on a hemocytometer. Cells were then plated atop thalamic neurons at 9 DIV at a concentration of 100,000 microglia/well. Following 24 hours, co-cultures were fixed in 4% PFA and 4% sucrose for 15 minutes at room temperature, washed in PBS, and mounted onto microscope slides in Fluoromount G + DAPI (SouthernBiotech). Imaging was performed on a Zeiss FV1000 confocal microscope as described above. Primary antibodies used include those listed above as well as mouse anti-PSD-95 (in-house), chicken anti-MAP2 (Lifespan Biosciences, LS-C61805), and rabbit anti-GFAP (Abcam, ab4674).

Statistical Analysis

N values for all experiments described in the main figures.

Figure 1. TEM analysis and Golgi staining

Eighty total boutons (40 with spines, 40 without spines) per condition were reconstructed and quantified in (E) and (F). (G), $n = 38$ (WT) and $n = 27$ (KO) bulbous spines. (H), $n = 68$ boutons per condition. (I), $n = 15$ WT and 15 KO neurons analyzed across 3 mice per condition. (J), $n = 35$ WT and 31 KO neurons analyzed across 3 mice per condition.

Figure 2. Golgi analysis of spines

(D), $n = 40$ (Cre-) and 29 (Cre+) neurons analyzed from 3 mice per condition. (E), $n = 30$ neurons across 3 mice per condition. (F), averaged spine lengths from $n = 39$ (Cre-) and 30 (Cre+) neurons from 3 mice per condition. (G), Averaged spine lengths from $n = 29$ neurons each from 3 mice per condition. In panels (I) - (O), $n = 30$ (P10, P90, and P27 LDR), 36 (P20), and 28 (P27 NR) neurons across 3 mice per condition.

Figure 3. Molecular analysis of TWEAK expression

(B) and (G): n = 3 mice per condition; (D), n = 8 (0) and 10 (8) mice per condition. (E), n = 54 (0) and 62 (8) microglia from 3 mice per condition. (F), n = 28 (0) and 44 (8) relay neurons from 3 mice per condition.

Figure 4. Engulfment assays and spine analysis

(B), n = 20 (P7 WT), 17 (P7 KO), 28 (P27 WT), and 37 (P27 KO) microglia from 4 mice per condition. (D), n = 42 (WT) and 30 (KO) microglia from 4 mice per condition. (F), n = 5 (P7 WT), 4 (P7 KO), 6 (P27 WT), and 8 (P27 KO) mice per condition. (G), n = 5 (P7 WT), 7 (P7 KO), 6 (P27 WT), and 9 (P27 KO) mice per condition. (J), conditions left to right, n = 25, 24, 24, 29, 38, and 37 neurons from 3 mice per condition.

Figure 5. Spine analyses and TWEAK localization

(B), conditions left to right, n = 40, 29, 29, and 31 neurons from 3 mice per condition. (D) – (I), n = 30 neurons across 3 mice per condition.

Figure 6. Microglia-neuron co-cultures

For (C) – (F), n = 51 (control), 47 (WT), and 45 (KO) neurons from 3 independent rounds of co-cultures. For (G) and (H), n = 104 and 117 bulbous spines, WT and KO respectively, pooled across 3 independent rounds of co-cultures. For (I) and (J), N = 174 (WT) and 228 (KO) thin spines across three separate bioreplicates.

Figure 7. FISH-IF analysis of microglial TWEAK and proximal synapses.

(C) – (G), n = 60 total microglia, 28 low TWEAK expressers and 32 high TWEAK expressers.

Blinding

All experiments were performed with investigators blinded to the conditions at each stage. For GridTape analysis, Fn14 KO and WT littermates were identified by S.R. via genotyping, and the mouse numbers were given to L.C. L.C. perfusion-fixed brains, microdissected out the dLGNs, and provided both sections to the Electron Microscopy Core for further processing (see above). The core stained then provided the coded sections to J.M-S. and W-C.L., who sectioned and aligned the datasets. L.C. then performed quantifications and reconstructions. Following all data analysis, L.C. was unblinded to the conditions.

For spine analysis, L.C. performed Golgi staining of comparable samples in parallel and blinded S.R. to the sample identities of brains upon freezing. S.R. then sectioned, stained, and quantified spines prior to being unblinded by L.C. For microglial engulfment assay, Sholl assay, and eye-specific segregation, S.R. blinded L.C. to the samples and L.C. performed imaging in parallel. L.C. then further blinded the samples for analysis by S.R. All other analyses, including RNAscope and immunoprecipitation, were similarly blinded.

Statistical tests and software used

Data processing, figure generation, and statistical analyses were performed in Graphpad Prism 7 and 8. For the comparison of parameters including only 2 conditions, Student's t test (unpaired) was used. For the comparison of more than 2 conditions within the same graph, One-way ANOVAs were performed, with post hoc comparison of all conditions to one another by Tukey's test. For comparison of microglial levels of TWEAK to other cell types, One-way ANOVA with Dunnett's test was used. Cumulative frequency distributions were statistically analyzed using Kolmogorov-Smirnov test.